

RESEARCH ARTICLE

Diagnosing Polyparasitism in a High-Prevalence Setting in Beira, Mozambique: Detection of Intestinal Parasites in Fecal Samples by Microscopy and Real-Time PCR

Lynn Meurs^{1,2}, Anton M. Polderman¹, Natalie V. S. Vinkeles Melchers^{1#a}, Eric A. T. Brienen¹, Jaco J. Verweij^{1#b}, Bernhard Groosjohan³, Felisberto Mendes³, Manito Mechendura³, Dagmar H. Hepp^{1#c}, Marijke C. C. Langenberg¹, Rosanne Edelenbosch^{1#d}, Katja Polman², Lisette van Lieshout^{1*}

1 Department of Parasitology, Leiden University Medical Center, Leiden, The Netherlands, **2** Department of Biomedical Sciences, Institute of Tropical Medicine, Antwerp, Belgium, **3** Faculty of Health Science, Catholic University of Mozambique, Beira, Mozambique


#a Current address: Department of Public Health, University Medical Center Rotterdam (ErasmusMC), Rotterdam, The Netherlands

#b Current address: Laboratory for Medical Microbiology and Immunology, St. Elisabeth Hospital, Tilburg, The Netherlands

#c Current address: Neurology Department, VU University Medical Center, Amsterdam, The Netherlands

#d Current address: Rathenau Institute, The Hague, The Netherlands

* E.A.van_Lieshout@lumc.nl



OPEN ACCESS

Citation: Meurs L, Polderman AM, Vinkeles Melchers NVS, Brienen EAT, Verweij JJ, Groosjohan B, et al. (2017) Diagnosing Polyparasitism in a High-Prevalence Setting in Beira, Mozambique: Detection of Intestinal Parasites in Fecal Samples by Microscopy and Real-Time PCR. *PLoS Negl Trop Dis* 11(1): e0005310. doi:10.1371/journal.pntd.0005310

Editor: William Evan Secor, Centers for Disease Control and Prevention, UNITED STATES

Received: June 26, 2016

Accepted: January 9, 2017

Published: January 23, 2017

Copyright: © 2017 Meurs et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: Leiden University Fund (LUF; www.luf.nl/en), the Jo Keur Fund (<https://www.lumc.nl/sub/9500/att/131107035332416.pdf>) and the LUSTRA scholarship (<http://hum.leiden.edu/internationalisation/outgoing-students/outgoing-exchange-students/news/lustra-scholarship>).

Abstract

Background

Many different intestinal parasite species can co-occur in the same population. However, classic diagnostic tools can only frame a particular group of intestinal parasite species. Hence, one or two tests do not suffice to provide a complete picture of infecting parasite species in a given population. The present study investigated intestinal parasitic infections in Beira, Mozambique, i.e. in the informal settlement of Inhamudima. Diagnostic accuracy of five classical microscopy techniques and real-time PCR for the detection of a broad spectrum of parasites was compared.

Methodology/Principal Findings

A cross-sectional population-based survey was performed. One stool sample per participant (n = 303) was examined by direct smear, formal-ether concentration (FEC), Kato smear, Baermann method, coproculture and real-time PCR. We found that virtually all people (96%) harbored at least one helminth, and that almost half (49%) harbored three helminths or more. Remarkably, *Strongyloides stercoralis* infections were widespread with a prevalence of 48%, and *Ancylostoma* spp. prevalence was higher than that of *Necator americanus* (25% versus 15%), the hookworm species that is often assumed to prevail in East-Africa. Among the microscopic techniques, FEC was able to detect the broadest spectrum of parasite species. However, FEC also missed a considerable number of infections, notably

[applications.html](#)) were awarded to MCCL, DHH and RE. The PCR analyses at Leiden University Medical Center were partly supported by the Prof. Dr. P.C. Flu-Foundation. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

S. stercoralis, *Schistosoma mansoni* and *G. intestinalis*. PCR outperformed microscopy in terms of sensitivity and range of parasite species detected.

Conclusions/Significance

We showed intestinal parasites—especially helminths—to be omnipresent in Inhamudima, Beira. However, it is a challenge to achieve high diagnostic sensitivity for all species. Classical techniques such as FEC are useful for the detection of some intestinal helminth species, but they lack sensitivity for other parasite species. PCR can detect intestinal parasites more accurately but is generally not feasible in resource-poor settings, at least not in peripheral labs. Hence, there is a need for a more field-friendly, sensitive approach for on-the-spot diagnosis of parasitic infections.

Author Summary

In populations living in adverse conditions due to poverty, a wide variety of intestinal parasite infections can be observed. These infections are usually diagnosed by stool microscopy, but can be easily missed if the procedures used are inaccurate or performed in a suboptimal way. In the present study, we investigated the prevalence of intestinal parasite infections in an informal settlement in Beira, Mozambique. We also compared the diagnostic performance of five different microscopic techniques and real-time PCR for the detection of DNA of five helminth and five protozoa species. For this purpose, a single stool sample was collected from members of 63 households, evenly distributed within the settlement. Parasitic infections were found to be highly abundant in the 303 examined samples. Virtually all individuals were found to be infected with at least one helminth species and a majority of the stools showed multiple parasites. None of the classical microscopic techniques was suitable to detect all different pathogenic species. Hence, the outcome of several microscopy procedures had to be combined to get a complete picture. We found PCR to be the most accurate diagnostic approach, even in this hyper-endemic setting. However, DNA detection is a high-tech technology, generally not applicable in resource-poor settings.

Introduction

Intestinal parasitic infections are among the most prevalent infections in humans in low- and middle-income countries. They can be largely categorized into two groups, i.e. helminthic and protozoan infections. Intestinal parasitic infections can cause significant morbidity. Especially children—who are generally more prone to heavy worm burdens—suffer from the sequelae of intestinal parasitic infections, such as diarrhea, malabsorption and anemia [1;2].

The most important intestinal helminths, both in terms of abundance and disease burden, are soil-transmitted helminths (STHs) such as hookworms, *Ascaris lumbricoides*, and *Trichuris trichiura* [3]. It is estimated that STHs infect more than two billion people or more than a third of the world's population [4]. Also, the *Schistosoma* spp. blood flukes are of great public health importance, with more than 250 million people infected worldwide [5;6], and an estimated global disease burden of 4.0 million disability-adjusted life years (DALYs) [7].

In comparison to helminth infections, less is known about intestinal protozoan infections. They have been associated with persistent diarrhea in developing countries [8–10], and can cause severe morbidity, especially in immunocompromised individuals [11]. Hundreds of millions of people may be affected by intestinal protozoa annually [12;13]. Yet, there are no reliable estimates of the global burden of disease [14–16]. This lack of knowledge is due to the fact that intestinal protozoa are difficult to diagnose. Also, some diagnostic techniques cannot distinguish pathogenic from non-pathogenic species (i.e. *Entamoeba histolytica* versus the other *Entamoeba* spp.). For some species there is no consensus on their pathogenicity (e.g. *Blastocystis*), while for others, disease only develops in certain infected individuals but not in all (e.g. *Giardia intestinalis*). Loss of microscopic skills in many clinical laboratories and the general lack of awareness on protozoan infections further add to these difficulties.

The diagnosis of intestinal parasites typically relies on the microscopic detection of egg, larval, trophozoite, cyst, and/or oocyst life stages in human feces samples [17;18]. The sensitivity of stool microscopy is generally low, and for a reliable diagnosis it is important to choose the appropriate microscopic technique [19]. For example, relatively simple techniques such as the direct smear are known to detect high *A. lumbricoides* loads while underestimating the presence of other helminths such as *Schistosoma mansoni* [20]. Ideally, the technique with the highest diagnostic accuracy for the parasite of interest should be selected. In practice however, this is difficult to achieve since many different parasite species may occur in a given population, or even in a single individual, and resources are generally limited in countries where most of these infections are endemic, so not all appropriate microscopic techniques can be used. In the past decade, alternative diagnostic procedures have become available, such as the detection of parasite DNA in stool samples using real-time PCR [21]. The disadvantage of PCR, however, is that—in contrast to microscopy—it needs a high-tech laboratory, which is even more of a challenge for diagnostic laboratories within endemic countries.

Relatively little is known about the distribution of intestinal parasites in Mozambique [22;23]. The present study was initiated because a local hospital noticed many cases of diarrhea in one of the informal settlements (‘bairro’) in Beira, Mozambique. Given the sanitary conditions in the study area, intestinal parasites were suspected to be the cause of these complaints. However, diagnostic methods that were being used in the hospital at that time were not adequate to detect these infections. Hence, the aim of this study was 1) to investigate which intestinal parasite species are most common in this area, and 2) to compare diagnostic accuracy between different microscopic techniques and real-time PCR for these intestinal parasitic infections. Five commonly used microscopic techniques were applied and evaluated, i.e. direct smear, formal-ether concentration (FEC), Kato smear, Baermann method, and coproculture, for the detection of a uniquely broad spectrum of intestinal parasites: from intestinal helminths like *Strongyloides stercoralis*, *Ancylostoma* spp., *Necator americanus*, *A. lumbricoides*, *T. trichiura* and *Schistosoma* spp. blood flukes, to pathogenic intestinal protozoa such as *G. intestinalis*, *E. histolytica*, the coccidium *Cystoisospora belli* and the microsporidia *Enterocytozoon bieneusi* and *Encephalitozoon* spp. Microscopy and real-time PCR results were compared to one another and to composite reference standards (CRSs).

Materials and Methods

Ethics statement

Approval to perform the study was obtained from the Beira Committee of Medical Ethics, Mozambique and the study proposal was filed by the Committee of Medical Ethics of the Leiden University Medical Centre (reference number CI5.151/NV/ib). Prior to the study, written informed consent was obtained from the head of participating households. Individuals who

were infected according to microscopy were offered treatment following standard clinical practice at the local hospital. Samples were anonymized for further data analysis.

Study population

The study was performed in Inhamudima (E34.86°, S19.84°), an informal settlement in the city of Beira, Mozambique, and was conducted on request of the local hospital and faculty of medicine. The area of Inhamudima is frequently flooded and is not connected to a sewage system. The rainy season lasts from October to March. The study was performed between June and August 2007.

A geographical map of this area was prepared and households and roads were annotated. In order to obtain a random and geographically evenly distributed sample of households and a logistically feasible sample size, a grid with 75 x 75 meter quadrants was superimposed on this map and the household that was closest to each of the intersections was selected. In this way, all participants of in total 63 households were approached to participate. In the field, these houses were located using handheld GPS devices. Infants (younger than one year) and people who did not provide sufficient fecal material for all procedures were excluded from the study. Fig 1 shows that 303 out of the 399 individuals that had given informed consent provided sufficient fecal material for inclusion into the study (i.e. participation rate of 76%).

Initially, urine samples were also collected for detection of *Schistosoma haematobium* (by urine filtration on one 10ml urine sample). Because of the relatively low numbers of *S. haematobium* cases however, collection of urine samples was stopped to focus on the diagnosis of intestinal parasites.




Fig 1. Flow diagram of the selection of the study population and diagnostic procedures.

doi:10.1371/journal.pntd.0005310.g001

Microscopy

Fecal samples were collected from all participating household members on a door-to-door basis, 0–18h after production of the samples, and examined in Beira within 24h after collection. Three well trained microscopists performed the laboratory procedures, and on average not more than eight stool samples were processed per day to ensure high quality microscopic results. Multiple approaches were used for the detection of cysts and oocysts of the protozoa, and eggs and larvae of the helminths (Table 1). Microscopic techniques included direct smear, FEC, Kato thick smear, Baermann method, and charcoal plate coproculture [17].

Table 1. Diagnostic methods performed and definitions of composite reference standards.

Parasite species	Direct smear	FEC	Kato smear	Baermann	Coproculture	PCR
STHs	<i>Strongyloides stercoralis</i>	+	+	-	+	+
	Hookworm	+	+	+	-	+ ^a
	<i>Ascaris lumbricoides</i>	+	+	+	-	+
	<i>Trichuris trichiura</i>	+	+	+	-	-
<i>Schistosoma mansoni</i> ^b	+	+	+ ^c	-	-	+ ^d
Pathogenic protozoa	<i>Giardia intestinalis</i>	+	+	-	-	+
	<i>Entamoeba histolytica</i> complex ^e	+	+	-	-	+
	<i>Cryptosporidium parvum</i> / <i>C. hominis</i> ^f	-	-	-	-	+ ^g
	<i>Enterocytozoon bieneusi</i>	-	-	-	-	+ ^g
	<i>Encephalitozoon</i> spp.	-	-	-	-	+ ^g
	<i>Cystoisospora belli</i> ^f	+	+	-	-	-
	<i>Cyclospora cayetanensis</i> ^f	-	-	-	-	-
Non-pathogenic protozoa	<i>Entamoeba coli</i>	+	+	-	-	-
	<i>Entamoeba hartmanni</i>	+	+	-	-	-
	<i>Iodamoeba bütschlii</i>	+	+	-	-	-
	<i>Endolimax nana</i>	+	+	-	-	-
	<i>Chilomastix mesnili</i>	+	+	-	-	-
	<i>Blastocystis</i> ^h	+	+	-	-	-
	<i>Dientamoeba fragilis</i> ^h	-	-	-	-	-

FEC, formal-ether concentration

For each parasite species, the composite reference standard (CRS) is based on the combined findings of the test(s) indicated with a plus sign (+). The minus sign (-) indicates the tests that were not applicable for a given parasite species. The latter tests were not included in the respective CRSs.

^a In contrast to the microscopic techniques, the PCR differentiates between the hookworm species *Ancylostoma* spp. and *Necator americanus*.

^b In addition to the microscopic methods mentioned, glycerin sedimentation was carried out [17]. However, this was stopped after the first 100 samples because it was very labor-intensive and did not detect any additional *S. mansoni* cases.

^c The time interval between preparation of the Kato smear and the examination of the slides was 30–60 minutes and therefore suboptimal for the detection of *S. mansoni*.

^d The PCR detects DNA of both *S. mansoni* and *S. haematobium*.

^e *Entamoeba histolytica* cannot be differentiated microscopically from *E. dispar* and the other *Entamoeba* spp. of the *E. histolytica* complex: *E. bangladeshi*, *E. ecuadoriensis*, *E. moshkovskii*, and *E. nutalli* [24–26]. The PCR was specific for *E. histolytica*. Consequently, this parasite was not included in the analysis on diagnostic accuracy.

^f The microscopic methods used—direct smear and FEC—are known to be inadequate for these protozoan infections. Hence, the modified Ziehl-Neelsen staining [17] was performed but stopped after the first 201 samples because it was labor-intensive and *Cryptosporidium* was detected in three samples only, while *C. belli* and *C. cayetanensis* were absent in this subsample. Because of incompleteness the modified Ziehl-Neelsen staining, data was not included in the CRS.

^g These protozoa cannot be detected by any of the microscopic techniques used on all samples, and were therefore not included in the analysis on diagnostic accuracy.

^h As yet, the classification of *Blastocystis* and *D. fragilis* as pathogenic or non-pathogenic species remains controversial [27;28].

doi:10.1371/journal.pntd.0005310.t001

For the direct smear, ~2mg of feces was mixed with normal saline on a microscopy slide and examined for helminth eggs. Another ~2mg of feces was mixed with a drop of iodine and examined for protozoan cysts [17]. For FEC, the fecal parasite concentrator (FPC, Evergreen) was used. One gram of fecal material was thoroughly mixed with 8 ml of 10% formalin. An FPC strainer with 15 ml tube was attached to the tube containing this mixture. After having filtered the suspension into the empty tube, 3 ml of ether was added to the filtrate. This mixture was then shaken vigorously for 1 minute and centrifuged at 500 x g for 2 minutes. A thick, unstained wet mount of the sediment was used for the detection of helminth eggs and larvae. For protozoan cysts, a thin, iodine-stained wet mount of the sediment was used.

The Kato smear—also known as Kato-Katz smear—consisted of a single slide of fecal material [18;29;30]. A 25 mg template was placed on the microscopy slide and filled with sieved (~300 µm pore size) fecal material. Upon removal of the template, the sample was covered with a cellophane slip soaked with glycerol and water (1:1). The sample was flattened by pressing it onto an even surface, and examined 30–60 minutes after preparation.

For the Baermann method, fecal material (~4g) was placed on a layer of 2 hydrophilic gauze bandages. The gauze was folded into a pouch by attaching the four perforated corners of the gauze to a stick. Subsequently, the pouch was placed in a 50 ml tube filled with tap water for 3h in such a way that the pouch lightly touched the water. Most of the water was decanted and the remaining sediment was left to stand for 2 hours before being examined for nematode larvae.

For coproculture, the classical charcoal culture procedure was used [31]. Approximately 2g of fecal material was homogenized, mixed 1:1 with vermiculite, and placed on a filter paper on a plastic platform in a petri dish. Tap water was added to wet the filter paper and the petri dish was covered. After incubation at room temperature for 7 days, the water was collected in a tube and left standing for 2h. The sediment was examined for nematode larvae.

For the Baermann method as well as for coproculture, two microscopy slides were prepared, each with 100µl of the sediment. A drop of iodine was added if moving larvae were detected, enabling identification and quantification of the larvae.

Real-time PCR

In Beira, an aliquot (~1g) of each stool sample was sieved and mixed with 3 volumes of 96% ethanol for preservation and shipment to Leiden, the Netherlands [32]. Here, the samples were stored at -20°C until detection and quantification of parasite DNA loads by real-time PCR. DNA isolation, amplification and detection were performed blinded to previous microscopic results.

For DNA isolation, 250µl of feces suspension was centrifuged and the pellet was washed with phosphate-buffered saline, resuspended in 200µl of 2% polyvinylpyrrolidone (Sigma) and heated for 10 minutes at 100°C [32;33]. After sodiumdodecyl sulfate-proteinase K treatment (2h at 55°C), DNA was isolated using QIAamp Spin Columns/Mini Kit (Qiagen, Germany). In each sample, a fixed amount of Phocine Herpes Virus 1 was included within the isolation lysis buffer as an internal control [34].

In total, 10 PCR targets were included and 5 µl DNA was used in each real-time PCR. Amplification generally comprised of 15' at 95°C followed by 50 cycles of 15" at 95°C, 30" at 60°C, and 30" at 72°C. Parasite-specific primers and probes were used for amplification of sequences, according to previously published protocols. Hookworm DNA (*Ancylostoma* spp. and *N. americanus*) was detected by one multiplex PCR described by Verweij et al. [35]. *Ascaris lumbricoides* [36] and *S. stercoralis* [37] DNA was detected in separate PCRs instead of in a multiplex format combined with other helminth targets. Schistosomal DNA was detected in

an additional PCR as described by Obeng et al. [38–41]. Protozoa (*E. histolytica*, *G. intestinalis*, and *C. parvum/C. hominis*) DNA was detected by multiplex HGC-PCR [42]. Microsporidial (*E. bienersi* and *Encephalitozoon* spp.) DNA was detected in another multiplex PCR described by Verweij et al. [43].

Negative and positive control samples were included in each PCR run. The PCR output from this system consisted of a cycle-threshold (Ct) value, representing the amplification cycle in which the level of fluorescent signal exceeded the background fluorescence. Hence, low Ct values correspond to high parasite-specific DNA loads in the sample tested, and vice versa. The maximum Ct value was set at 50 indicating that DNA was not detected in the sample after 50 cycles of amplification. The Ct values of the internal Phocine Herpes Virus 1 control were within the expected range (Ct value between 30 and 33) for all samples, indicating that there was no evidence of inhibition of amplification in any of these samples.

Analysis

IBM SPSS 22.0 (IBM Corp.) and Microsoft Excel 14.0 (Microsoft Corp.) were used for statistical analyses. GraphPad Prism 5 (GraphPad Software, Inc.) was used to prepare graphs.

There is no gold standard for the detection of individual intestinal parasite species. Although microscopic techniques are known to lack sensitivity, they are, just as the PCR, supposed to be 100% specific [44]. We therefore combined the results of several diagnostic methods into a composite reference standard (CRS) [45;46]. The CRS was defined in such a way that it was negative if none of the diagnostic methods detected the parasite of interest, and positive if one or more methods detected the parasite. Table 1 shows how the CRS was defined for the different parasite species.

Infection prevalence was based on the CRS unless stated otherwise, and 95% Wald confidence intervals were calculated for this parameter. For sensitivities of the different diagnostic methods, the Wilson score method without continuity correction was used to calculate 95% confidence intervals [47]. Differences between test sensitivities were considered statistically significant if there was no overlap of their confidence intervals. The independent samples Mann-Whitney U test was used to determine whether differences in Ct values between microscopy-positives and -negatives were statistically significant.

Results

Characteristics of the study population

The study population (n = 303) consisted of 144 (48%) males and 159 (52%) females with a median age of 17 years (range 1 to 72). These people were derived from 63 households. Per household 1 to 11 subjects participated (median of 4 subjects).

Infection prevalence

STH infections were widespread with a prevalence of 93%, 56%, 48% and 38% for *T. trichiura*, *A. lumbricoides*, *S. stercoralis* and hookworm, respectively (Fig 2A). PCR indicated that *Ancylostoma* spp. was the most abundant hookworm: 25% of the population (75/303) harbored *Ancylostoma* spp. while *N. americanus* was detected in 15% (46/303) of the population. Mixed *Ancylostoma* spp. and *N. americanus* infections were observed in 5% (15/303) of the population. The prevalence of *S. mansoni* was 10.9%.

Within the population of Inhamudima, 96% (292/303) of individuals were found to harbor at least one of the following helminths in their stool sample: *S. stercoralis*, *Ancylostoma* spp., *N. americanus*, *A. lumbricoides*, *T. trichiura*, *S. mansoni*, and 49% (147/303) of the total




Fig 2. Prevalence of intestinal parasitic infections in the study population according to different diagnostic methods. Whiskers indicate 95% confidence intervals of the observed prevalence. Percentages are based on observations in 303 individuals. (A) Prevalence of helminth infections. *Strongyloides stercoralis* infection was not determined (ND) in Kato smears, hookworm was not determined by the Baermann method, while *A. lumbricoides*, *T. trichiura*, and *S. mansoni* were not determined by the Baermann method or coproculture. *Trichuris trichiura* was not determined by PCR either, and the composite reference standard (CRS) for this infection was consequently based on microscopic results only. (B) Prevalence of intestinal protozoan infections. Feces were examined by both microscopy and PCR for *G. intestinalis*, and *E. histolytica* complex spp. (one observation was missing for PCR, and consequently for the CRS). Only PCR data was used for *C. parvum/C. hominis* (one observation missing) and for, *E. bienuesi* and *Encephalitozoon* spp. (two observations missing). * While microscopy cannot differentiate between the pathogenic species *Entamoeba histolytica* and the nonpathogenic species of the *E. histolytica* complex, PCR is specific for the pathogenic species (*E. histolytica*).

doi:10.1371/journal.pntd.0005310.g002

population harbored three or more different helminth species (Fig 3). Up to five helminth infections were found in one individual.

The prevalence of infections with pathogenic protozoa is shown in Fig 2B. The pathogenic protozoan *Cystoisospora belli* was not detected in this population. *Giardia intestinalis* was the




Fig 3. Number of helminth species found per person. Prevalence of infection is based on the composite reference standard for *S. stercoralis*, *A. lumbricoides*, *T. trichiura*, and *S. mansoni*, and on PCR for hookworm—*Ancylostoma* spp. and *N. americanus* (n = 303).

doi:10.1371/journal.pntd.0005310.g003

most common pathogenic protozoan with a prevalence of 37% (113/302). Microsporidia were found in 9% (28/301) and mainly consisted of *E. bienersi* infections (27/28 cases). *Cryptosporidium parvum/C. hominis* was found by PCR in 2% (6/302) of the study population. All showed a Ct value higher than 30 (median Ct 37.6) and none of these 6 overlapped with the 3 samples positive in the Ziehl-Neelsen staining (legend f, Table 1). The prevalence of *E. histolytica* complex spp. infections was 10% (30/303) based on microscopy. However, PCR showed that only 10% (3/30) of these infections involved *E. histolytica*, i.e. the pathogenic species. Combining the findings of pathogenic protozoa with the detection of helminths, 98% (294/300) of the tested inhabitants of Inhamudima were found to be infected with at least one intestinal parasite species.

The prevalence of non-pathogenic protozoa varied between 4% and 34% for the different species, with a prevalence of 34% (102/303) for *Entamoeba coli*, 26% (80/303) for *Endolimax nana*, 21% (63/302) for *Blastocystis*, 9% (26/303) for *Entamoeba hartmanni*, 8% (24/303) for *Chilomastix mesnili* and a prevalence of 4% (13/303) for *Iodamoeba bütschlii*.

Comparison of microscopic techniques

Diagnostic sensitivity was estimated for the different microscopic techniques and for each of the parasite species (Fig 4). FEC and Kato smear had the highest sensitivities for the detection of each of the helminths, except for *S. stercoralis*. The direct smear was inferior to these two methods for the detection of *S. stercoralis*, hookworm, *T. trichiura* and *S. mansoni*. Similarly, the direct smear was inferior to FEC for the detection of *G. intestinalis*. The Baermann method and coproculture showed the highest sensitivities for *S. stercoralis* (48% and 57%, respectively), while the direct smear and FEC showed equally low sensitivities (18% and 25%, respectively). The sensitivity of coproculture for the detection of *S. stercoralis* and hookworm (57% and 77%, respectively) tended to be higher than the sensitivity of the other microscopic techniques (up to 48% and 69%, respectively). Moreover, the sensitivity of helminth detection increased upon combination of multiple microscopic methods.




Fig 4. Sensitivities of the different diagnostic methods for the detection of intestinal parasitic infections. Whiskers indicate 95% confidence intervals of observed sensitivities (n = 303). *Strongyloides stercoralis* infection was not determined (ND) in Kato smears, hookworm was not determined by the Baermann method, while *A. lumbricoides*, *T. trichiura*, and *S. mansoni* were not determined by the Baermann method or coproculture. *Trichuris trichiura* was not determined by PCR, and for this species the sensitivity was therefore based on microscopic results only. *Giardia intestinalis* was not determined by Kato smear, the Baermann method or coproculture (one observation missing).

doi:10.1371/journal.pntd.0005310.g004

Microscopy versus PCR

The sensitivity of PCR for the detection of each of the parasite species tested was higher than that of any of the microscopic techniques applied (Fig 4). This difference was statistically significant for all microscopic methods used for *S. stercoralis*, hookworm, *S. mansoni*, and *G. intestinalis*, and for the direct smear for the detection of *A. lumbricoides*. For some species, the sensitivity of the best microscopic technique was manifold lower than that of PCR. For example, the estimated sensitivity for the detection of *S. mansoni* was 12% for Kato smear versus 97% for the detection of *Schistosoma* DNA in feces via PCR. For *G. intestinalis*, sensitivity of FEC was 21% while that of PCR was 99%. In addition, in only one of the three PCR-positive *E. histolytica* samples, *E. histolytica* complex spp. cysts could be detected by microscopy (FEC).

Fig 5 shows that, for each parasite species, PCR-positive but microscopy-negative samples had significantly lower DNA loads (i.e. higher Ct values) than PCR-positive samples that were also microscopy-positive. Although observed less frequently, some microscopy-positive samples could not be confirmed by PCR. In most of these samples, only few parasites were detected by microscopy (see Table 2).

Discussion

The present study was initiated because a local hospital noticed many cases of diarrhea in Inhamudima. Given the adverse living conditions in the study area, intestinal parasites were suspected to be the cause of these complaints. However, diagnostic methods that were being used in the hospital at that time were not adequate. Clinical records thus far suggested that there were only low numbers of intestinal parasite infections in the area. The first aim of the present study was to obtain a comprehensive overview of the intestinal helminth and protozoan infections that occur in the informal settlement of Inhamudima in Beira.




Fig 5. Ct values in PCR-positives: microscopy-negative versus positive samples. Total number of PCR-positives per species is indicated between brackets. '-' indicates microscopy-negative and '+' microscopy-positive samples. Horizontal lines indicate median Ct values. Differences in Ct values between microscopy-positive and -negative samples were all significant $p \leq 0.007$, ^a microscopy cannot differentiate the two hookworm species.

doi:10.1371/journal.pntd.0005310.g005

The major strength of the present study is the unprecedentedly large panel of different diagnostic techniques used [48]. This diagnostic rigor resulted in high quality data regarding a uniquely wide spectrum of intestinal parasite species, including helminths as well as protozoa. Prevalences observed here are thus likely to approach the true prevalence of individual parasitic infections as well as co-infections [49], in contrast to other prevalence studies where generally only one microscopic technique is used. This approach led to several important observations concerning the presence of intestinal parasite infections in the target population, such as the remarkably high levels of *S. stercoralis*, and high parasite infection levels in general, as well as a high degree of polyparasitism.

According to Schär et al., only 44% of African countries have prevalence data on *S. stercoralis*, and a majority of these studies is based on inadequate diagnostic techniques [50]. In Mozambique, so far, only Mandomando et al. reported on *S. stercoralis*. A prevalence of 1.1% was reported in under-five-year-olds with diarrhea referred to a district hospital in Southern Mozambique, but only direct smear was used which is ineffective for the detection of this parasite [51]. As a result, *S. stercoralis* prevalences were assumed to be low, i.e. 6.2% prevalence in Mozambique [50]. Our results (prevalence of 48%) indicate that this may be a gross

Table 2. Discordance between microscopy and PCR in microscopy-positives.

Parasite species	Number of microscopy-positive cases being PCR-negative (number of parasites observed per technique)						Total number of microscopy-positives
	Direct smear	FEC	Kato smear	Baermann	Coproculture	Total	
<i>S. stercoralis</i>	1 (1)	0	NA	7 ^a	5 (6–141) ^a	12	100
Hookworm	0	1 (2) ^b	3 (1–47) ^b	NA	7 (2–1000) ^b	9	98
<i>A. lumbricoides</i>	2 (1)	5 (1–4)	0	NA	NA	7	161
<i>S. mansoni</i>	0	0	1 (1) ^c	NA	NA	1	5
<i>G. intestinalis</i> ^d	0	1 (1–10)	NA	NA	NA	1	24

FEC, Formal-ether concentration; NA, Not applicable

^a One case was detected by coproculture as well as the Baermann method. In addition to stool, Baermann medium and coproculture medium were analyzed by PCR in the 7 Baermann-positives and 3/5 coproculture-positives, respectively. These samples also tested negative.

^b One case was detected by FEC, Kato smear as well as coproculture.

^c In addition, 3 *S. haematobium* eggs were observed

^d Parasite numbers were determined semi-quantitatively.

doi:10.1371/journal.pntd.0005310.t002

underestimation of the nationwide prevalence and once more illustrate how often strongyloidiasis is overlooked due to poor diagnostics.

Another important observation was that *Ancylostoma* spp. was more prevalent than *N. americanus* (25% versus 15%, respectively). This is in line with the studies described by Goldsmid [52], as well as a more recent study in pre-school children living in a rural area in Malawi where *Ancylostoma* spp. prevailed over *N. americanus* [53]. Despite of these reports, it is often assumed that only *N. americanus* is endemic in East Africa [54].

High levels of intestinal parasitic infections as observed in the present study are not exceptional in urban informal settlements. More than 20 years ago, it was already predicted that ongoing urbanization in the developing world would increase levels of intestinal parasites in areas like Inhamudima [55]. In such adverse living conditions, widespread environmental contamination is likely to occur for parasites such as *A. lumbricoides*, *T. trichiura*, and *G. intestinalis* [55]. The present observations confirm this idea as these three parasites ranked in the top four of most prevalent parasites in Inhamudima, with a prevalence as high as 93% for *T. trichiura*. Augusto et al. (2009) also noted a particularly high STH prevalence in (peri-)urban areas of Mozambique (including Beira, although these were not explicitly reported), and these prevalences were higher than those reported in older studies [22]. The authors hypothesized that the increase of STH infections in Mozambican informal settlements was due to rural-urban migration, low levels of socioeconomic development, and overcrowding together with the deterioration of water supply and sewage systems after independence [22].

In addition to determining infection levels of intestinal parasites in the inhabitants of Inhamudima, this study aimed to compare diagnostic accuracy between five microscopic techniques that are commonly used for the detection of intestinal parasites. The common microscopic techniques direct smear, FEC, Kato smear, Baermann method, and coproculture were applied and compared for the detection of parasitic helminths as well as protozoan infections. The classical broad-spectrum techniques FEC and Kato smear proved useful for the detection of intestinal helminths: hookworm, *T. trichiura* and *A. lumbricoides*. The Kato smear may be more informative than FEC as it is a quantitative technique, and in accordance with a recent meta-analysis by Nikolay et al., it tended to detect more STH infections—more hookworm, *T. trichiura*, and *A. lumbricoides*—than FEC [44]. On the other hand, the Kato smear is not suitable for the detection of pathogenic intestinal protozoa. In this respect, the two techniques are complementary. In order to also detect *S. stercoralis*, coproculture and/or the Baermann method should be added to the diagnostic work-up. Consistent with a recent meta-analysis by Campo Polanco et al. [56], coproculture tended to have a higher sensitivity for the detection of *S. stercoralis* than the Baermann method. We demonstrated that coproculture has the additional advantage that it has a higher sensitivity for the detection of hookworm than the other techniques, while the Baermann method is suited for the detection of *S. stercoralis* only. This was also reported by Knopp et al. [48]. The disadvantage of coproculture however, is that it takes a week to obtain test results while the Baermann method takes several hours, and that it is more difficult to perform under field conditions [48]. In addition, coproculture harbors the occupational hazard of infection by L3 larvae. In concordance with earlier reports [44], the direct smear was inferior to all the other diagnostic tests, for each of the endemic intestinal parasite species. This confirms that the direct smear is not at all suitable for the diagnosis of intestinal parasite infections. In practice however, it is widely used in endemic areas, in both research and clinical settings—including the Beira hospital—resulting in gross underestimations of the spread of intestinal parasites.

We aimed to compare the diagnostic accuracy of the different microscopic techniques to that of real-time PCR. The detection of parasite-specific DNA had a higher sensitivity than microscopy for *S. stercoralis*, hookworm, *Schistosoma* spp., and *G. intestinalis*, even when

results from the five microscopic techniques were combined. These findings are in line with earlier studies, despite differences in endemicity and geographic locations [44]. For example, Arndt et al. showed the same helminth PCRs that were used in the present study (in a multiplex format [36]) to be more sensitive for the detection of *S. stercoralis*, hookworm and *Schistosoma* spp. than the combination of direct smear, FEC, and Kato smear [57]. Also Easton et al. recently reported that real-time PCR outperformed Kato-Katz for the detection of soil-transmitted helminths and has the advantage of demonstrating parasite species which are not diagnosed by Kato-Katz such as *G. intestinalis*, *E. histolytica* and *S. stercoralis* [58]. Earlier findings from our group comparing Kato smear and PCR in Kenyan schoolchildren showed that PCR outperformed the Kato smear for the detection of *N. americanus* as well as *S. mansoni*. Even when three consecutive stool samples were analyzed using Kato smear, while PCR was performed on only one stool sample, PCR detected more cases [41;59].

When focusing on the diagnosis of *Schistosoma* only, previous studies found PCR to be more sensitive than microscopy, [41], also when using other *Schistosoma*-specific targets [60–64]. It should be noted that *S. haematobium* was detected in our study populations. This parasite was seen in six out of 95 examined urine samples (6%). PCR results indicated that *Schistosoma* DNA was absent in stool samples from people with *S. haematobium* eggs in urine but without *S. mansoni* infection according to microscopy. In accordance with the Kenyan study [41], *S. haematobium* co-endemicity is thus unlikely to explain the relatively low sensitivity of microscopy as compared to PCR for the detection of *S. mansoni*. Yet, we cannot entirely rule out that “leakage” of *S. haematobium* DNA may have resulted in occasional false-positive PCRs.

To our knowledge, only one study reported a lower diagnostic accuracy of PCR relative to microscopy. Knopp et al. showed real-time PCR for the detection of hookworm and of *S. stercoralis* to be less sensitive than microscopy when applied on stool samples collected in a low-endemicity area in Tanzania [65]. Although this study used the same PCR procedures as in the present study, this inconsistency could still be explained by minor technical differences in the performance of the DNA extraction or the general PCR set-up, resulting in a reduced test efficiency which becomes more obvious when infection levels are already low. Hence, standardization of laboratory procedures and the implementation of external quality assessment schemes are warranted.

It is widely recognized that PCR could be particularly useful for the detection of intestinal parasites in low transmission areas and in post-control settings [48;58]. Here, we show that PCR can also have added value in high transmission areas, particularly in polyparasitized populations. It not only has a higher sensitivity than microscopy for all the intestinal parasites investigated in this study, but it also has the advantage that it can differentiate between morphologically identical species where microscopy cannot. This is important because infections caused by species passing morphologically indistinguishable eggs, such as the hookworms, can have different epidemiology. Moreover, it also detects infections such as microsporidia which are extremely difficult to diagnose by conventional light microscopy. Although it was not yet possible to detect *T. trichiura* by PCR at the time of study, currently available multiplex PCRs do include this helminth. PCR may thus be a useful tool for evaluation of public health interventions, for quality control of microscopy procedures as well as for research on the distribution of intestinal parasitic infections in different endemic settings [46;57]. However, it should be noted that PCR also has disadvantages. PCR consumables are expensive, and it requires high-tech equipment which is often not available in laboratories in affected regions. Transfer of samples to other, better equipped laboratories, may prolong the turnaround time of PCR results. Secondly, the clinical relevance of submicroscopic intestinal parasitic infections is not yet fully understood, and therefore remains a matter of debate [16;57;66]. It has also been

proposed that residual DNA may persist and be detected by PCR after parasite clearance [67], while others have showed fast DNA clearance after treatment [68;69]. More research is needed to clarify these issues.

Conclusion

We demonstrate that intestinal helminth and protozoan infections and co-infections are widespread in Inhamudima, Beira. We showed that classical techniques such as FEC are useful for the detection of some intestinal helminths such as hookworm, *T. trichiura* and *A. lumbricoides*. However, they lack the sensitivity to reliably characterize the wide range of intestinal parasites that may coexist in a population or individual. PCR can detect intestinal parasites more accurately but in most endemic areas it is not (yet) possible to perform this technique, at least not in the more peripheral laboratories. So, until a more field-friendly approach becomes available, infection levels of intestinal parasites—and polyparasitism—are best approximated by combining multiple and relatively simple microscopic techniques.

Supporting Information

S1 Checklist.

(DOCX)

S1 Database.

(XLSX)

Acknowledgments

We gratefully thank the population of Inhamudima, Beira, Mozambique for participation in this study, Idelson for washing materials, and Sandra for translating. Also, we would like to thank the anonymous reviewer who so generously shared his/her encyclopedic parasitology knowledge.

Author Contributions

Conceptualization: AMP.

Data curation: LvL LM.

Formal analysis: LM.

Funding acquisition: AMP LvL MCCL RE DHH.

Investigation: EATB DHH MCCL RE MM FM.

Methodology: AMP LvL JJV.

Project administration: AMP LvL BG.

Resources: BG.

Supervision: AMP LvL.

Visualization: LM AMP.

Writing – original draft: LM.

Writing – review & editing: LM JJV AMP NVSVM KP LvL DHH MCCL RE.

References

1. Guerrant RL, Oria RB, Moore SR, Oria MO, Lima AA. Malnutrition as an enteric infectious disease with long-term effects on child development. *Nutr Rev* 2008 Sep; 66(9):487–505. doi: [10.1111/j.1753-4887.2008.00082.x](https://doi.org/10.1111/j.1753-4887.2008.00082.x) PMID: [18752473](https://pubmed.ncbi.nlm.nih.gov/18752473/)
2. Hall A, Hewitt G, Tuffrey V, de Silva N. A review and meta-analysis of the impact of intestinal worms on child growth and nutrition. *Matern Child Nutr* 2008 Apr; 4 Suppl 1:118–236.
3. Hotez PJ, Kamath A. Neglected tropical diseases in sub-saharan Africa: review of their prevalence, distribution, and disease burden. *PLoS Negl Trop Dis* 2009; 3(8):e412. doi: [10.1371/journal.pntd.0000412](https://doi.org/10.1371/journal.pntd.0000412) PMID: [19707588](https://pubmed.ncbi.nlm.nih.gov/19707588/)
4. Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted helminth infections: ascariasis, trichuriasis, and hookworm. *Lancet* 2006 May 6; 367(9521):1521–32. doi: [10.1016/S0140-6736\(06\)68653-4](https://doi.org/10.1016/S0140-6736(06)68653-4) PMID: [16679166](https://pubmed.ncbi.nlm.nih.gov/16679166/)
5. Colley DG, Bustinduy AL, Secor WE, King CH. Human schistosomiasis. *Lancet* 2014 Mar 31.
6. Gryseels B, Polman K, Clerinx J, Kestens L. Human schistosomiasis. *Lancet* 2006 Sep 23; 368(9541):1106–18. doi: [10.1016/S0140-6736\(06\)69440-3](https://doi.org/10.1016/S0140-6736(06)69440-3) PMID: [16997665](https://pubmed.ncbi.nlm.nih.gov/16997665/)
7. WHO. Health statistics and information systems: Estimates for 2000–2012: Disease burden. 2016. 6-5-2016.
8. Becker SL, Vogt J, Knopp S, Panning M, Warhurst DC, Polman K, et al. Persistent digestive disorders in the tropics: causative infectious pathogens and reference diagnostic tests. *BMC Infect Dis* 2013; 13:37. doi: [10.1186/1471-2334-13-37](https://doi.org/10.1186/1471-2334-13-37) PMID: [23347408](https://pubmed.ncbi.nlm.nih.gov/23347408/)
9. Pawlowski SW, Warren CA, Guerrant R. Diagnosis and treatment of acute or persistent diarrhea. *Gastroenterology* 2009 May; 136(6):1874–86. doi: [10.1053/j.gastro.2009.02.072](https://doi.org/10.1053/j.gastro.2009.02.072) PMID: [19457416](https://pubmed.ncbi.nlm.nih.gov/19457416/)
10. Polman K, Becker SL, Alirof E, Bhatta NK, Bhattarai NR, Bottieau E, et al. Diagnosis of neglected tropical diseases among patients with persistent digestive disorders (diarrhoea and/or abdominal pain >/= 14 days): A multi-country, prospective, non-experimental case-control study. *BMC Infect Dis* 2015; 15(1):338.
11. Stark D, Barratt JL, van Hal S, Marriott D, Harkness J, Ellis JT. Clinical significance of enteric protozoa in the immunosuppressed human population. *Clin Microbiol Rev* 2009 Oct; 22(4):634–50. doi: [10.1128/CMR.00017-09](https://doi.org/10.1128/CMR.00017-09) PMID: [19822892](https://pubmed.ncbi.nlm.nih.gov/19822892/)
12. Lane S, Lloyd D. Current trends in research into the waterborne parasite *Giardia*. *Crit Rev Microbiol* 2002; 28(2):123–47. doi: [10.1080/1040-840291046713](https://doi.org/10.1080/1040-840291046713) PMID: [12109771](https://pubmed.ncbi.nlm.nih.gov/12109771/)
13. Checkley W, White AC Jr., Jaganath D, Arrowood MJ, Chalmers RM, Chen XM, et al. A review of the global burden, novel diagnostics, therapeutics, and vaccine targets for *Cryptosporidium*. *Lancet Infect Dis* 2015 Jan; 15(1):85–94. doi: [10.1016/S1473-3099\(14\)70772-8](https://doi.org/10.1016/S1473-3099(14)70772-8) PMID: [25278220](https://pubmed.ncbi.nlm.nih.gov/25278220/)
14. Stanley SL Jr. Amoebiasis. *Lancet* 2003 Mar 22; 361(9362):1025–34. doi: [10.1016/S0140-6736\(03\)12830-9](https://doi.org/10.1016/S0140-6736(03)12830-9) PMID: [12660071](https://pubmed.ncbi.nlm.nih.gov/12660071/)
15. Turkeltaub JA, McCarty TR III, Hotez PJ. The intestinal protozoa: emerging impact on global health and development. *Curr Opin Gastroenterol* 2015 Jan; 31(1):38–44. doi: [10.1097/MOG.0000000000000135](https://doi.org/10.1097/MOG.0000000000000135) PMID: [25394233](https://pubmed.ncbi.nlm.nih.gov/25394233/)
16. Efunshile MA, Ngwu BA, Kurtzhals JA, Sahar S, Konig B, Stensvold CR. Molecular Detection of the Carriage Rate of Four Intestinal Protozoa with Real-Time Polymerase Chain Reaction: Possible Overdiagnosis of *Entamoeba histolytica* in Nigeria. *Am J Trop Med Hyg* 2015 Aug 5; 93(2):257–62. doi: [10.4269/ajtmh.14-0781](https://doi.org/10.4269/ajtmh.14-0781) PMID: [26101274](https://pubmed.ncbi.nlm.nih.gov/26101274/)
17. Garcia LS. Diagnostic medical parasitology. 2007. LWW.
18. WHO. Basic laboratory methods in medical parasitology. Geneva: World Health Organization; 1991.
19. van Lieshout L, Roestenberg M. Clinical consequences of new diagnostic tools for intestinal parasites. *Clin Microbiol Infect* 2015 Apr 2.
20. van Lieshout L, Yazdanbakhsh M. Landscape of neglected tropical diseases: getting it right. *Lancet Infect Dis* 2013 Jun; 13(6):469–70. doi: [10.1016/S1473-3099\(13\)70094-X](https://doi.org/10.1016/S1473-3099(13)70094-X) PMID: [23562237](https://pubmed.ncbi.nlm.nih.gov/23562237/)
21. Verweij JJ, Stensvold CR. Molecular testing for clinical diagnosis and epidemiological investigations of intestinal parasitic infections. *Clin Microbiol Rev* 2014 Apr; 27(2):371–418. doi: [10.1128/CMR.00122-13](https://doi.org/10.1128/CMR.00122-13) PMID: [24696439](https://pubmed.ncbi.nlm.nih.gov/24696439/)
22. Augusto G, Nala R, Casmo V, Sabonete A, Mapaco L, Monteiro J. Geographic distribution and prevalence of schistosomiasis and soil-transmitted helminths among schoolchildren in Mozambique. *Am J Trop Med Hyg* 2009 Nov; 81(5):799–803. doi: [10.4269/ajtmh.2009.08-0344](https://doi.org/10.4269/ajtmh.2009.08-0344) PMID: [19861614](https://pubmed.ncbi.nlm.nih.gov/19861614/)
23. Clavero AO, Verdu ME, Peman J, Dario R, Gobernado M. Human intestinal infection due to coccidia in Mozambique: two cases. *Acta Trop* 1999 Jan 15; 72(1):25–9. PMID: [9924958](https://pubmed.ncbi.nlm.nih.gov/9924958/)

24. Parija SC, Mandal J, Ponnambath DK. Laboratory methods of identification of *Entamoeba histolytica* and its differentiation from look-alike *Entamoeba* spp. *Trop Parasitol* 2014 Jul; 4(2):90–5. doi: [10.4103/2229-5070.138535](https://doi.org/10.4103/2229-5070.138535) PMID: [25250228](https://pubmed.ncbi.nlm.nih.gov/25250228/)
25. Clark CG, Kaffashian F, Tawari B, Windsor JJ, Twigg-Flesner A, Davies-Morel MC, et al. New insights into the phylogeny of *Entamoeba* species provided by analysis of four new small-subunit rRNA genes. *Int J Syst Evol Microbiol* 2006 Sep; 56(Pt 9):2235–9. doi: [10.1099/ij.s.0.64208-0](https://doi.org/10.1099/ij.s.0.64208-0) PMID: [16957127](https://pubmed.ncbi.nlm.nih.gov/16957127/)
26. Levecke B, Dorny P, Vercammen F, Visser LG, Van EM, Vercruyssen J, et al. Transmission of *Entamoeba nuttalli* and *Trichuris trichiura* from Nonhuman Primates to Humans. *Emerg Infect Dis* 2015 Oct; 21(10):1871–2. doi: [10.3201/eid2110.141456](https://doi.org/10.3201/eid2110.141456) PMID: [26402309](https://pubmed.ncbi.nlm.nih.gov/26402309/)
27. Clark CG, van der Giezen M, Alfellani MA, Stensvold CR. Recent developments in *Blastocystis* research. *Adv Parasitol* 2013; 82:1–32. doi: [10.1016/B978-0-12-407706-5.00001-0](https://doi.org/10.1016/B978-0-12-407706-5.00001-0) PMID: [23548084](https://pubmed.ncbi.nlm.nih.gov/23548084/)
28. Garcia LS. *Dientamoeba fragilis*, One of the Neglected Intestinal Protozoa. *J Clin Microbiol* 2016 Sep; 54(9):2243–50. doi: [10.1128/JCM.00400-16](https://doi.org/10.1128/JCM.00400-16) PMID: [27053676](https://pubmed.ncbi.nlm.nih.gov/27053676/)
29. Katz N, Chaves A, Pellegrino J. A simple device for quantitative stool thick-smear technique in *Schistosomiasis mansoni*. *Rev Inst Med Trop Sao Paulo* 1972 Nov; 14(6):397–400. PMID: [4675644](https://pubmed.ncbi.nlm.nih.gov/4675644/)
30. Polderman AM, Mpamila K, Manshande JP, Bouwhuis-Hoogerwerf ML. Methodology and interpretation of parasitological surveillance of intestinal schistosomiasis in Maniema, Kivu Province, Zaire. *Ann Soc Belg Med Trop* 1985 Sep; 65(3):243–9. PMID: [3935059](https://pubmed.ncbi.nlm.nih.gov/3935059/)
31. Polderman AM, Eberhard M, Baeta S, Gasser RB, van Lieshout L, Magnussen P, et al. Chapter 3—The Rise and Fall of Human Oesophagostomiasis. In: *Advances in Parasitology*. Volume 71 ed. Academic Press; 2010. p. 93–155.
32. Verweij JJ, Oostvogel F, Brienen EA, Nang-Beifubah A, Ziem J, Polderman AM. Prevalence of *Entamoeba histolytica* and *Entamoeba dispar* in northern Ghana. *Trop Med Int Health* 2003 Dec; 8(12):1153–6. PMID: [14641852](https://pubmed.ncbi.nlm.nih.gov/14641852/)
33. Verweij JJ, Polderman AM, Wimmenhove MC, Gasser RB. PCR assay for the specific amplification of *Oesophagostomum bifurcum* DNA from human faeces. *Int J Parasitol* 2000 Feb; 30(2):137–42. PMID: [10704596](https://pubmed.ncbi.nlm.nih.gov/10704596/)
34. Niesters HG. Clinical virology in real time. *J Clin Virol* 2002 Dec; 25 Suppl 3:S3–12.
35. Verweij JJ, Brienen EA, Ziem J, Yelifari L, Polderman AM, van Lieshout L. Simultaneous detection and quantification of *Ancylostoma duodenale*, *Necator americanus*, and *Oesophagostomum bifurcum* in fecal samples using multiplex real-time PCR. *Am J Trop Med Hyg* 2007 Oct; 77(4):685–90. PMID: [17978072](https://pubmed.ncbi.nlm.nih.gov/17978072/)
36. Wiria AE, Prasetyani MA, Hamid F, Wammes LJ, Lell B, Ariawan I, et al. Does treatment of intestinal helminth infections influence malaria? Background and methodology of a longitudinal study of clinical, parasitological and immunological parameters in Nangapanda, Flores, Indonesia (ImmunoSPIN Study). *BMC Infect Dis* 2010; 10:77. doi: [10.1186/1471-2334-10-77](https://doi.org/10.1186/1471-2334-10-77) PMID: [20338054](https://pubmed.ncbi.nlm.nih.gov/20338054/)
37. Verweij JJ, Canales M, Polman K, Ziem J, Brienen EA, Polderman AM, et al. Molecular diagnosis of *Strongyloides stercoralis* in faecal samples using real-time PCR. *Trans R Soc Trop Med Hyg* 2009 Apr; 103(4):342–6. doi: [10.1016/j.trstmh.2008.12.001](https://doi.org/10.1016/j.trstmh.2008.12.001) PMID: [19195671](https://pubmed.ncbi.nlm.nih.gov/19195671/)
38. Obeng BB, Aryeetey YA, de Dood CJ, Amoah AS, Larbi IA, Deelder AM, et al. Application of a circulating-cathodic-antigen (CCA) strip test and real-time PCR, in comparison with microscopy, for the detection of *Schistosoma haematobium* in urine samples from Ghana. *Ann Trop Med Parasitol* 2008 Oct; 102(7):625–33. doi: [10.1179/136485908X337490](https://doi.org/10.1179/136485908X337490) PMID: [18817603](https://pubmed.ncbi.nlm.nih.gov/18817603/)
39. Pillay P, Taylor M, Zulu SG, Gundersen SG, Verweij JJ, Hoekstra P, et al. Real-Time Polymerase Chain Reaction for Detection of *Schistosoma* DNA in Small-Volume Urine Samples Reflects Focal Distribution of Urogenital Schistosomiasis in Primary School Girls in KwaZulu Natal, South Africa. *Am J Trop Med Hyg* 2014 Mar; 90(3):546–52. doi: [10.4269/ajtmh.13-0406](https://doi.org/10.4269/ajtmh.13-0406) PMID: [24470560](https://pubmed.ncbi.nlm.nih.gov/24470560/)
40. Vinkeles Melchers NV, van Dam GJ, Shaproski D, Kahama AI, Brienen EA, Vennervald BJ, et al. Diagnostic performance of *Schistosoma* real-time PCR in urine samples from Kenyan children infected with *Schistosoma haematobium*: day-to-day variation and follow-up after praziquantel treatment. *PLoS Negl Trop Dis* 2014 Apr; 8(4):e2807. doi: [10.1371/journal.pntd.0002807](https://doi.org/10.1371/journal.pntd.0002807) PMID: [24743389](https://pubmed.ncbi.nlm.nih.gov/24743389/)
41. Meurs L, Brienen E, Mbow M, Ochola EA, Mboup S, Karanja DM, et al. Is PCR the Next Reference Standard for the Diagnosis of *Schistosoma* in Stool? A Comparison with Microscopy in Senegal and Kenya. *PLoS Negl Trop Dis* 2015 Jul; 9(7):e0003959. doi: [10.1371/journal.pntd.0003959](https://doi.org/10.1371/journal.pntd.0003959) PMID: [26217948](https://pubmed.ncbi.nlm.nih.gov/26217948/)
42. Bruijnesteijn van Coppenraet LE, Wallinga JA, Ruijs GJ, Bruins MJ, Verweij JJ. Parasitological diagnosis combining an internally controlled real-time PCR assay for the detection of four protozoa in stool samples with a testing algorithm for microscopy. *Clin Microbiol Infect* 2009 Sep; 15(9):869–74. doi: [10.1111/j.1469-0691.2009.02894.x](https://doi.org/10.1111/j.1469-0691.2009.02894.x) PMID: [19624500](https://pubmed.ncbi.nlm.nih.gov/19624500/)

43. Verweij JJ, ten Hove R, Brienen EA, van Lieshout L. Multiplex detection of *Enterocytozoon bieneusi* and *Encephalitozoon* spp. in fecal samples using real-time PCR. *Diagn Microbiol Infect Dis* 2007 Feb; 57(2):163–7. doi: [10.1016/j.diagmicrobio.2006.08.009](https://doi.org/10.1016/j.diagmicrobio.2006.08.009) PMID: [17020796](https://pubmed.ncbi.nlm.nih.gov/17020796/)
44. Nikolay B, Brooker SJ, Pullan RL. Sensitivity of diagnostic tests for human soil-transmitted helminth infections: a meta-analysis in the absence of a true gold standard. *Int J Parasitol* 2014 Oct 1; 44(11):765–74. doi: [10.1016/j.ijpara.2014.05.009](https://doi.org/10.1016/j.ijpara.2014.05.009) PMID: [24992655](https://pubmed.ncbi.nlm.nih.gov/24992655/)
45. Alonzo TA, Pepe MS. Using a combination of reference tests to assess the accuracy of a new diagnostic test. *Stat Med* 1999 Nov 30; 18(22):2987–3003. PMID: [10544302](https://pubmed.ncbi.nlm.nih.gov/10544302/)
46. Glinz D, Silue KD, Knopp S, Lohourignon LK, Yao KP, Steinmann P, et al. Comparing diagnostic accuracy of Kato-Katz, Koga agar plate, ether-concentration, and FLOTAC for *Schistosoma mansoni* and soil-transmitted helminths. *PLoS Negl Trop Dis* 2010; 4(7):e754. doi: [10.1371/journal.pntd.0000754](https://doi.org/10.1371/journal.pntd.0000754) PMID: [20651931](https://pubmed.ncbi.nlm.nih.gov/20651931/)
47. Newcombe RG. Two-sided confidence intervals for the single proportion: comparison of seven methods. *Stat Med* 1998 Apr 30; 17(8):857–72. PMID: [9595616](https://pubmed.ncbi.nlm.nih.gov/9595616/)
48. Knopp S, Mgeni AF, Khamis IS, Steinmann P, Stothard JR, Rollinson D, et al. Diagnosis of soil-transmitted helminths in the era of preventive chemotherapy: effect of multiple stool sampling and use of different diagnostic techniques. *PLoS Negl Trop Dis* 2008; 2(11):e331. doi: [10.1371/journal.pntd.0000331](https://doi.org/10.1371/journal.pntd.0000331) PMID: [18982057](https://pubmed.ncbi.nlm.nih.gov/18982057/)
49. Booth M, Vounatsou P, N'goran EK, Tanner M, Utzinger J. The influence of sampling effort and the performance of the Kato-Katz technique in diagnosing *Schistosoma mansoni* and hookworm co-infections in rural Cote d'Ivoire. *Parasitology* 2003 Dec; 127(Pt 6):525–31. PMID: [14700188](https://pubmed.ncbi.nlm.nih.gov/14700188/)
50. Schar F, Trostorf U, Giardina F, Khieu V, Muth S, Marti H, et al. *Strongyloides stercoralis*: Global Distribution and Risk Factors. *PLoS Negl Trop Dis* 2013; 7(7):e2288. doi: [10.1371/journal.pntd.0002288](https://doi.org/10.1371/journal.pntd.0002288) PMID: [23875033](https://pubmed.ncbi.nlm.nih.gov/23875033/)
51. Mandomando IM, Macete EV, Ruiz J, Sanz S, Abacassamo F, Valles X, et al. Etiology of diarrhea in children younger than 5 years of age admitted in a rural hospital of southern Mozambique. *Am J Trop Med Hyg* 2007 Mar; 76(3):522–7. PMID: [17360878](https://pubmed.ncbi.nlm.nih.gov/17360878/)
52. Goldsmid JM. The African Hookworm Problem. Parasitic helminths and zoonoses in Africa. 1 ed. London: Unwin Hyman; 1991. p. 101–30.
53. Jonker FA, Calis JC, Phiri K, Brienen EA, Khoffi H, Brabin BJ, et al. Real-time PCR demonstrates *Ancylostoma duodenale* is a key factor in the etiology of severe anemia and iron deficiency in Malawian pre-school children. *PLoS Negl Trop Dis* 2012; 6(3):e1555. doi: [10.1371/journal.pntd.0001555](https://doi.org/10.1371/journal.pntd.0001555) PMID: [22514750](https://pubmed.ncbi.nlm.nih.gov/22514750/)
54. Palmer PES, Reeder MM. Ancylostomiasis (hookworm disease). In: The imaging of tropical diseases: With epidemiological, pathological and clinical correlation. 2nd ed. Heidelberg: Springer-Verlag; 2001. p. 51–65.
55. Crompton DW, Savioli L. Intestinal parasitic infections and urbanization. *Bull World Health Organ* 1993; 71(1):1–7. PMID: [8440028](https://pubmed.ncbi.nlm.nih.gov/8440028/)
56. Campo Polanco L, Gutierrez LA, Cardona Arias J. Diagnosis of *Strongyloides stercoralis* infection: meta-analysis on evaluation of conventional parasitological methods (1980–2013). *Rev Esp Salud Publica* 2014 Oct; 88(5):581–600. doi: [10.4321/S1135-57272014000500004](https://doi.org/10.4321/S1135-57272014000500004) PMID: [25327268](https://pubmed.ncbi.nlm.nih.gov/25327268/)
57. Arndt MB, John-Stewart G, Richardson BA, Singa B, van Lieshout L, Verweij JJ, et al. Impact of helminth diagnostic test performance on estimation of risk factors and outcomes in HIV-positive adults. *PLoS One* 2013; 8(12):e81915. doi: [10.1371/journal.pone.0081915](https://doi.org/10.1371/journal.pone.0081915) PMID: [24324729](https://pubmed.ncbi.nlm.nih.gov/24324729/)
58. Easton AV, Oliveira RG, O'Connell EM, Kepha S, Mwandawiro CS, Njenga SM, et al. Multi-parallel qPCR provides increased sensitivity and diagnostic breadth for gastrointestinal parasites of humans: field-based inferences on the impact of mass deworming. *Parasit Vectors* 2016; 9:38. doi: [10.1186/s13071-016-1314-y](https://doi.org/10.1186/s13071-016-1314-y) PMID: [26813411](https://pubmed.ncbi.nlm.nih.gov/26813411/)
59. van Mens SP, Aryeetey Y, Yazdanbakhsh M, van Lieshout L, Boakye D, Verweij JJ. Comparison of real-time PCR and Kato smear microscopy for the detection of hookworm infections in three consecutive faecal samples from schoolchildren in Ghana. *Trans R Soc Trop Med Hyg* 2013 Apr; 107(4):269–71. doi: [10.1093/trstmh/trs094](https://doi.org/10.1093/trstmh/trs094) PMID: [23426113](https://pubmed.ncbi.nlm.nih.gov/23426113/)
60. Allam AF, Kader O, Zaki A, Shehab AY, Farag HF. Assessing the marginal error in diagnosis and cure of *Schistosoma mansoni* in areas of low endemicity using Percoll and PCR techniques. *Trop Med Int Health* 2009 Mar; 14(3):316–21. doi: [10.1111/j.1365-3156.2009.02225.x](https://doi.org/10.1111/j.1365-3156.2009.02225.x) PMID: [19278527](https://pubmed.ncbi.nlm.nih.gov/19278527/)
61. Pontes LA, Dias-Neto E, Rabello A. Detection by polymerase chain reaction of *Schistosoma mansoni* DNA in human serum and feces. *Am J Trop Med Hyg* 2002 Feb; 66(2):157–62. PMID: [12135287](https://pubmed.ncbi.nlm.nih.gov/12135287/)

62. Pontes LA, Oliveira MC, Katz N, Dias-Neto E, Rabello A. Comparison of a polymerase chain reaction and the Kato-Katz technique for diagnosing infection with *Schistosoma mansoni*. *Am J Trop Med Hyg* 2003 Jun; 68(6):652–6. PMID: [12887022](#)
63. Gomes LI, Marques LH, Enk MJ, Coelho PM, Rabello A. Further evaluation of an updated PCR assay for the detection of *Schistosoma mansoni* DNA in human stool samples. *Mem Inst Oswaldo Cruz* 2009 Dec; 104(8):1194–6. PMID: [20140385](#)
64. Carvalho GC, Marques LH, Gomes LI, Rabello A, Ribeiro LC, Scopel KK, et al. Polymerase chain reaction for the evaluation of *Schistosoma mansoni* infection in two low endemicity areas of Minas Gerais, Brazil. *Mem Inst Oswaldo Cruz* 2012 Nov; 107(7):899–902. PMID: [23147146](#)
65. Knopp S, Salim N, Schindler T, Karagiannis Voules DA, Rothen J, Lweno O, et al. Diagnostic accuracy of Kato-Katz, FLOTAC, Baermann, and PCR methods for the detection of light-intensity hookworm and *Strongyloides stercoralis* infections in Tanzania. *Am J Trop Med Hyg* 2014 Mar; 90(3):535–45. doi: [10.4269/ajtmh.13-0268](#) PMID: [24445211](#)
66. Pierce KK, Kirkpatrick BD. Update on human infections caused by intestinal protozoa. *Curr Opin Gastroenterol* 2009 Jan; 25(1):12–7. PMID: [19119509](#)
67. Frickmann H, Schwarz NG, Rakotozandrindrainy R, May J, Hagen RM. PCR for enteric pathogens in high-prevalence settings. What does a positive signal tell us? *Infect Dis (Lond)* 2015 Jul; 47(7):491–8.
68. Mejia R, Vicuna Y, Broncano N, Sandoval C, Vaca M, Chico M, et al. A novel, multi-parallel, real-time polymerase chain reaction approach for eight gastrointestinal parasites provides improved diagnostic capabilities to resource-limited at-risk populations. *Am J Trop Med Hyg* 2013 Jun; 88(6):1041–7. doi: [10.4269/ajtmh.12-0726](#) PMID: [23509117](#)
69. van den Bijlaardt W, Overvest IT, Buiting AG, Verweij JJ. Rapid clearance of *Giardia lamblia* DNA from the gut after successful treatment. *Clin Microbiol Infect* 2014 Nov; 20(11):O972–O974. doi: [10.1111/1469-0691.12626](#) PMID: [24655130](#)