

Barcoding hybrids: heterogeneous distribution of *Schistosoma haematobium* × *Schistosoma bovis* hybrids across the Senegal River Basin

NELE A. M. BOON^{1,2}, FREDERIK VAN DEN BROECK³, DJIBY FAYE⁴, FILIP A. M. VOLCKAERT¹, SOULEYMANE MBOUP⁵, KATJA POLMAN² and TINE HUYSE^{6*}

¹Laboratory of Biodiversity and Evolutionary Genomics, Biology, University of Leuven, Ch. Deberiotstraat 32, B-3000 Leuven, Belgium

²Unit of Medical Helminthology, Department of Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155, B-2000 Antwerp, Belgium

³Unit of Molecular Parasitology, Unit of Veterinary Protozoology, Department of Biomedical Sciences, Institute of Tropical Medicine, Nationalestraat 155, B-2000 Antwerp, Belgium

⁴Sante Plus, 9756 Sacré Coeur 3, BP 11294 Dakar, Sénégal

⁵Laboratoire de Bactériologie, Virologie du CHNU Aristide Le Dantec, Dakar, Sénégal

⁶Department of Biology, Royal Museum for Central Africa, Leuvensesteenweg 13, B-3080 Tervuren, Belgium

(Received 7 October 2017; revised 9 February 2018; accepted 12 February 2018; first published online 18 April 2018)

ABSTRACT

Hybridization events between *Schistosoma* species (Digenea, Platyhelminthes) are reported with increasing frequency, largely due to improved access to molecular tools. Nevertheless, little is known about the distribution and frequency of hybrid schistosomes in nature. Screening for hybrids on a large scale is complicated by the need for nuclear and mitochondrial sequence information, precluding a ‘simple’ barcoding approach. Here we aimed to determine and understand the spatiotemporal distribution of *Schistosoma haematobium* × *Schistosoma bovis* hybrids in the Senegal River Basin. From ten villages, distributed over the four main water basins, we genotyped a total of 1236 schistosome larvae collected from human urine samples using a partial mitochondrial *cox1* fragment; a subset of 268 parasites was also genotyped using ITS rDNA. Hybrid schistosomes were unevenly distributed, with substantially higher numbers in villages bordering Lac de Guiers than in villages from the Lampsar River and the Middle Valley of the Senegal River. The frequency of hybrids per village was not linked with the prevalence of urinary schistosomiasis in that village. However, we did find a significant positive association between the frequency of hybrids per village and the prevalence of *Schistosoma mansoni*. We discuss the potential consequences of adopting a barcoding approach when studying hybrids in nature.

Key words: Hybridization, *Schistosoma haematobium*, *Schistosoma bovis*, barcoding, Senegal, ITS rDNA, hybrids.

INTRODUCTION

Schistosomiasis is a major poverty-related disease, affecting more than 200 million people in developing tropical countries. The genus *Schistosoma* comprises in total 26 species, all of which are infecting mammals and among which eight are infecting humans (Brant and Loker, 2013). The major burden of human infection resides in sub-Saharan Africa where more than 90% of cases are registered; 112 million people are estimated to be infected with *Schistosoma haematobium* and 54 million with *Schistosoma mansoni* (Brindley and Hotez, 2013). In addition, in Africa, three *Schistosoma* species are found with a significant impact on livestock: *Schistosoma mattheei*, *Schistosoma bovis* and *Schistosoma curassoni* (Leger and Webster, 2016).

* Corresponding author: Tine Huyse, Department of Biology, Royal Museum for Central Africa, Leuvensesteenweg 13, B-3080 Tervuren, Belgium. E-mail: Tine.Huyse@africamuseum.be

A DNA barcoding approach was used to study the genetic variation and population structure of *S. haematobium* (Webster *et al.* 2012) and *S. mansoni* (Morgan *et al.* 2005; Webster *et al.* 2013a) on a large geographical scale. Surprisingly low sequence variation was found within mitochondrial genes of *S. haematobium* (Webster *et al.* 2012), which is in stark contrast with the high mtDNA diversity found in *S. mansoni* (Morgan *et al.* 2005; Webster *et al.* 2013a). Two main geographical groups were found in *S. haematobium*: one is found throughout the African mainland and the other is found exclusively at the Indian Ocean Islands and the neighbouring African coastal regions. A barcoding approach has also proven to be an ideal tool to study the diversity and interrelationships within the freshwater snails in the genus of *Bulinus* (Kane *et al.* 2008), of which some species act as an intermediate host for *S. haematobium* and *S. bovis*.

Schistosoma worms have a complex life cycle. Their transmission depends on both an intermediate

snail host and a final vertebrate host. Schistosomes are the only genus of Trematoda with separate sexes; both males and females are required for sexual reproduction (Southgate *et al.* 1998; Loker and Brant, 2006). Mating of adults takes place at the hepatic portal system and is required for migration towards the egg-laying site and maturation of female worms (Southgate *et al.* 1998). Although most worms show species-specific mating preferences, heterospecific mating occurs, even when a same species mate is available (Norton *et al.* 2008). This may result in parthenogenesis or hybridization, depending on the schistosome species involved (Southgate *et al.* 1998). Upon mating, female worms release their eggs into the veins of the surrounding organs. Eggs migrate towards the bladder, liver and intestine, depending on the species and reach the environment through urine or feces.

The Senegal River Basin (SRB) represents a transmission hotspot for schistosomiasis. In 1988 the Diama Dam was constructed on the Senegal River and a year later the Mantali Dam was built on the Bafing River, an upstream tributary of the Senegal River in Mali. These dams provided fresh water for irrigation and new habitats for snails, triggering a huge outbreak of intestinal schistosomiasis in the late eighties. A few years following this outbreak the prevalence of *S. haematobium*, causing urinary schistosomiasis increased also (Picquet *et al.* 1996; Southgate *et al.* 2001). Today, *S. mansoni* and *S. haematobium* are co-endemic in the SRB and along the shores of Lac de Guiers, resulting in mixed *Schistosoma* infections in humans (Meurs *et al.* 2012; Abdellahi *et al.* 2016). The prevalence of *S. haematobium* still seems to increase while *S. mansoni* prevalence tends to decline (personal Observation: personal communication Elke Steinacher in 2014, poste de santé Catholique in Dagana). *Schistosoma bovis* and *S. curassoni* are also present in northern Senegal (Vercruyssen *et al.* 1984) but a recent study only found *S. bovis* in the SRB (Webster *et al.* 2013b). Both species cause intestinal schistosomiasis in ruminants.

The SRB (Fig. 1) was among the first places where hybridization between the human host species *S. haematobium* and the cattle host species *S. bovis* was observed (Huyse *et al.* 2009). This hybrid was first described by Brémond *et al.* (1993) in Niger. *Schistosoma haematobium* and *S. bovis* belong both to the *S. haematobium* species group. Several species pairs within this group are known to hybridize experimentally or in the wild (Tchuem Tchuente *et al.* 1996; Webster *et al.* 2006; Leger and Webster, 2016). Laboratory experiments have shown that hybrids may have an enhanced transmission potential. For example, F1 and F2 hybrids between *S. haematobium* and *S. guineensis* exhibited a greater infectivity for snail intermediate hosts than

the parental schistosome species (Webster and Southgate, 2003). In the SRB, several *Bulinus* species act as intermediate snail host for *S. haematobium*, the main intermediate snail host is *Bulinus globosus*. *Bulinus truncatus*, the most common bulinid snail in the SRB, is host of *S. bovis* (Diaw *et al.* 1998; Sène *et al.* 2004). If the *S. haematobium* × *S. bovis* hybrids show infectivity to both *B. globosus* and *B. truncatus*, this could induce an increased transmission of these hybrids and therefore lead to higher prevalences of urinary schistosomiasis. However, the exact intermediate snail host range of the hybrid species is unknown at present. The *S. haematobium* × *S. bovis* hybrid seems widespread and has also been found elsewhere in Senegal (Webster *et al.* 2013b), Corsica and Benin (Moné *et al.* 2015) and Mali (Soentjens *et al.* 2016). Also, hybridization between the two phylogenetically remotely related human species *S. haematobium* and *S. mansoni* has been found in the SRB (Huyse *et al.* 2013). Despite the common observation of hybrids in the SRB (Huyse *et al.* 2009; Webster *et al.* 2013b), little is known about their spatio-temporal distribution.

Our first aim was to determine and explain the spatiotemporal distribution of hybridization in the SRB. Hybrids were detected through barcoding, based on the genetic profile of a mitochondrial marker (partial cytochrome *c* oxidase subunit I (*cox1*)). We compared the frequency of *S. haematobium* × *S. bovis* hybrids in the SRB in ten villages, sampled between 2006 and 2010, along four regions in the SRB; the Delta and the Middle Valley of the Senegal River, the Lampsar River and Lac de Guiers (Fig. 1). Secondly, we aimed to understand the potential impact of *S. haematobium* × *S. bovis* hybrids on the local epidemiology of schistosomiasis. The recent increase in *S. haematobium* prevalence across the SRB might be linked with the presence of hybrids. Therefore the frequency of hybrids, per village, was related to the prevalence of urinary and intestinal schistosomiasis.

MATERIAL AND METHODS

Study area and data collection

We sampled parasite stadia (eggs and miracidia) for genetic analysis and parasitological information on the prevalence and infection intensity of schistosomiasis in ten villages located in four regions in Northern Senegal (Fig. 1). Data were collected during parasitological surveys in 2006, 2007, 2009 and 2010 (Table 1). From the village of Diatar (1993) material from adult worms was obtained from the Schistosomiasis Collection at the Natural History Museum (London, UK), where field-collected cercariae were passaged through mice (SCAN; Emery *et al.* 2012).




Fig. 1. Map with the sample locations in the Senegal River Basin including in the Middle Valley the villages of Diatar, Dioundiou, and Kodit; towards the coast, in the Senegal River Delta the village of Rhonne; in the vicinity of Lac de Guiers on the eastern shore the village of Mbane, on the western shore the villages of Ndieumeul and Diokhor Tack, and north of the lake the village of Pakh; in the Lampsar River, a tributary of the Senegal River, the villages Mbodienne and Pont Gendarme. For each region, the percentage of hybrids (i.e. parasites with a *S. bovis* *cox1* mtDNA profile) is indicated by a pie chart.

The prevalence of *S. haematobium* and *S. mansoni*

In March 2006 urine and stool samples were collected from 20 school-age children in each of the villages of Dioundiou, Kodit, Mbane, Ndieumeul, Mbodienne and Pont Gendarme (Table 1). In 2007 parasitological data were collected from 75 school-age children in Diatar, Dioundiou, Rhonne, Mbane, Ndieumeul and Mbodienne. In 2009 and 2010 parasitological data were collected from both children (older than 4) and adults at the villages of Pakh, Diokhor and Ndieumeul (Table 1). *Schistosoma haematobium* was detected by filtering 10 mL of urine on a Nucleopore filter (Isopore, USA). *Schistosoma mansoni* infection was diagnosed by a duplicate (two slides of 25 mg) Kato Katz (Katz et al. 1972) per stool sample. In the studies of 2006 and 2007, one urine and one stool sample were collected, in the studies of 2009 and 2010 two urine and two stool samples per person were collected. The prevalence of *Schistosoma* species in urine and stool are reported per village and per sampling year (Table 1). In the studies of 2006 and 2007 *S. haematobium* and *S. mansoni* prevalence is calculated as the proportion of infected school-aged children. In the studies of 2009 and 2010 prevalence is calculated as the proportion of infected individuals across all ages.

Molecular analysis of *S. haematobium* stadia

Schistosoma eggs from positive urine samples were isolated by sedimentation. Isolated eggs hatched through exposure to sunlight in Petri dishes with clean bottled water to hatch. Individual miracidia were pipetted onto FTA[®] Classic cards (Whatman) in a volume of 3 μ L. Eggs that did not hatch were stored on 100% ethanol. DNA extraction of eggs and miracidia was performed as described by Van den Broeck et al. (2011). DNA from adult worms was extracted using the Nucleospin[®] Tissue kit following manufacturer's instructions (Macherey-Nagel).

mtDNA genotyping. A rapid diagnostic 'multiplex' one-step polymerase chain reaction (RD-PCR) was performed to discriminate between *S. haematobium*, *S. bovis* and *S. mansoni* (Webster et al. 2010; Van den Broeck et al. 2011). A partial cytochrome *c* oxidase subunit I (*cox1*) mtDNA fragment of parasite DNA was amplified in a total reaction volume of 25 μ L consisting of 1 \times PCR buffer, 1.5 mM MgCl₂, 200 μ M of each dNTP, 0.4 μ M Asmit 1 (Eurogentec, forward primer TTTTTT TGGTCA-TCCTGAGGTGTAT, Bowles et al. 1992), 0.2 μ M of each reverse primer [*S. haematobium* (Sh.R.) TGATAATCAATGAC

Table 1. Overview of parasite barcoding and prevalence of schistosomiasis in Northern Senegal per geographical region, village and year of sampling; the ITS rDNA genotype and *cox1* haplotype of *Schistosoma haematobium* miracidia collected from urines; the prevalence of urinary and intestinal schistosomiasis.

Village	year	Barcoding parasites # human ind.	ITS rDNA		<i>cox1</i> mtDNA		Prevalence schistosomiasis		
			# parasites	Genotype ^a	# parasites	% <i>S.b</i>	# human ind.	urine	stool
Senegal River (Middle Valley)									
Diatar	1993 ⁺⁺	NA	9		11	0.0%	NA	NA	NA
	2007	3	13		27	0.0%	75	57%	8%
Dioundiou	2006	10 (pool ^b)	10		31	3.2%	22	95%	33%
	2007	3	3		71	2.8%	37	79%	6%
Kodit	2006	9 (pool ^b)	8	1 <i>S.h/S.b</i>	25	4.0%	20	100%	17%
<i>Total</i>			44		165	2.4%	154	83%	16%
Senegal River (Delta)									
Rhonne	2007	2	41	1 <i>S.m</i> 1 <i>S.h/S.m</i>	30	16.7%	75	73%	81%
Lac de Guiers									
Mbane	2006	Pool ^b	12		27	29.6%	20	89%	61%
	2007	4	0		42	26.2%	76	89%	38%
Ndieumeul	2006	5	7		40	37.5%	20	95%	94%
	2007	4	0		36	16.7%	75	93%	60%
<i>Total</i>			19		145	27.5%	191	92%	63%
Ndieumeul ^c	Winter 2009/2010	17	7		196	28.6%	99	66%	75%
Pakh ^c	2009	15	33	2 <i>S.h/S.b</i>	465	20.9%	530	58%	17%
Diokhor ^c	2009	1	0		23	73.9%	599	44%	55%
<i>Total</i>			42		661	25.7%	1419	56%	49%
Lampsar River									
Mbodienne	2006	3	23		28	7.1%	20	95%	60%
	2007	4	44		104	0.0%	82	63%	25%
Pont Gendarme	2006	3	21		29	3.5%	20	85%	35%
<i>Total</i>			88		161	1.9%	122	81%	40%

Ind, individuals; NA, not applicable, *S.h*, *S. haematobium*; *S.b*, *S. bovis*; *S.m*, *S. mansoni*.

^a Genotype: The ITS rDNA genotype is only reported when it displayed a hybrid or a *S. mansoni* profile. In all other cases, the ITS rDNA profile showed a *S. haematobium* genotype. The abbreviations *S.h /S.b* and *S.h /S.m* indicate that at the diagnostic sites two chromatogram peaks were visible after Sanger sequencing.

^b Pool: In 2006, *Schistosoma* eggs were isolated by sedimentation after pooling positive urine samples from several human individuals. For Mbane (2006) the exact number of human urines that was pooled is not known.

^c These studies cover the whole population of a village, including school-age children and adults.

⁺⁺ Adult worms.

CCTGCAATAA 543 bp (Webster *et al.* 2010), *S. bovis* (Sb.R) CACAGGATCAGACAAACGA GT ACC 306 bp (Webster *et al.* 2010), *S. mansoni* (Sm.R) TGCAGATAAAGCCACCCCTGTG 375 bp (Van den Broeck *et al.* 2011)], 1 unit *Taq* polymerase (Eurogentec), milli-Q water and 2 μ L of DNA. PCR parameters during thermal cycling were 3 min at 95 °C, followed by 40 cycles of 30 s at 94 °C, 1 min 30 s at 58 °C and 1 min 30 s at 72 °C followed by a final cycle at 72 °C for 7 min. PCR products were visualized on a 1% Midori Green agarose gel.

ITS rDNA genotyping. Samples were subjected to PCR amplification of the complete rDNA region (981 bp; primers ITS4: TCCTCCGCTTATTGATATGC and ITS5: GGAAGTAAAAGT-CGT AACAAG (Barber *et al.* 2000)). PCR amplification

was performed in a volume of 25 μ L containing 1 \times PCR buffer, 2 mM MgCl₂, 200 μ M of each dNTP, 2 mM of each primer, 1 unit *Taq* polymerase (Eurogentec), milli-Q water and 2 μ L of DNA. Thermal cycling started with 3 min at 95 °C, followed by 40 cycles of 30 s at 94 °C, 45 s at 54 °C and 1 min 30 s at 72 °C followed by a final cycle at 72 °C for 7 min. We brought 4 μ L of the PCR product on a 1% Midori Green agarose gel to visualize whether the PCR reaction was successful. PCR products of successful PCR reactions were further analysed by restriction fragment length polymorphism (RFLP) and visualized on a 1% Midori Green agarose gel (See Figures S1 and S2, supplementary information). Digestion of the ITS fragment with *MboI* (Fermentas) distinguishes *S. haematobium* from *S. bovis* and *S. mansoni*. *Schistosoma*

haematobium PCR products were digested in fragment sizes of (in bp) 280, 174, 130, 100, 85 and 36 bp; *S. bovis* in fragment sizes of 380, 174, 167, 130, 83 and 36 bp; for *S. mansoni* in sizes of 380, 176, 174, 130, 81, 16 and 9 bp. Samples showing a *S. bovis* or *S. mansoni MboI* pattern were subsequently digested with *AluI* (Fermentas), which differentiates between *S. bovis* (no cut; 981 bp) and *S. mansoni* (fragments of 545 and 421 bp). We used for each RFLP 8.8 μL of the same PCR product in a reaction of 10 μL (1 μL buffer, 0.2 μL enzyme) for digestion with the *MboI* and the *AluI* enzyme, respectively. To confirm the genotype based on restriction enzymes, the ITS rDNA region was sequenced for 19 randomly chosen parasite larvae by Sanger sequencing. The PCR products were purified using the E.Z.N.A Cycle Pure kit (Omega Bio-tek, Norcross, USA). Sanger sequencing was done using a Big Dye Chemistry Cycle Sequencing kit v3.1 in a 3130 Genetic Analyzer (Applied Biosystems). Sequences were assembled and manually edited using Geneious[®] v6.1.6. (<http://geneious.com>).

Hybrid definition

The hybrid definition used in this study builds on barcoding; hybrids are defined based on the genetic profile of a mitochondrial marker (partial cytochrome *c* oxidase subunit I (*cox1*)) and a nuclear marker (ITS rDNA). If discordance between both profiles was found (mito-nuclear discordance) the parasite was defined as ‘hybrid’. If both markers showed a *S. haematobium* profile, the parasite was defined as *S. haematobium*, although introgression of other nuclear *S. bovis* genes cannot be ruled out (Boon *et al.* 2017). Parasites with a *S. bovis cox1* profile that lacked an ITS rDNA profile were also defined as *S. haematobium* \times *S. bovis* hybrids, assuming that human infection by pure *S. bovis* parasites is absent or at most rare (Boissier *et al.* 2016). Hence, in this study, the frequency of hybrids is a synonym for the frequency of eggs or miracidia with a *S. bovis cox1* profile.

Data analyses

R v3.3.1-based packages (R Development Core Team, 2016) were used for all statistical analyses. The genetic profile of the mitochondrial *cox1* fragment and the nuclear ITS rDNA was reported for each parasite stadium (adult worms, eggs and miracidia). To test whether the frequency of hybrids differed among the four regions, a Kruskal–Wallis test (Kruskal and Wallis, 1952) was performed with the frequency of hybrids per village as dependent variable and region as grouping factor. Afterwards, pairwise comparisons among the four regions were performed using Dunn’s test (Dunn,

1964) with a Bonferroni correction for multiple comparisons. Both tests were performed with the R-package *dunn.test* (Dinno, 2016).

We explored with a principal component analysis (PCA) with R-package *ade4* (Dray and Dufour, 2007) the correlation between the frequency of hybrids per village, *S. haematobium* prevalence and *S. mansoni* prevalence. The prevalence of *S. haematobium* and *S. mansoni* was based on microscopy-based egg counts, and the frequency of hybrids was based on the molecular identification of eggs or miracidia. The statistical significance of the association between the frequency of hybrids per village and *S. mansoni* prevalence was investigated with a linear regression based on the percentage of hybrids per village (separated per year of sampling) as the dependent variable and *S. mansoni* prevalence as the independent variable.

RESULTS

Species identification of the mitochondrial cox1 and nuclear ITS marker

The number and percentage of parasites with a *Schistosoma bovis*, *S. haematobium* or *S. mansoni* profile for each genetic marker is presented in Table 2. The *cox1* fragment was successfully genotyped in 1236 parasite stadia; the ITS rDNA was successfully genotyped by RFLP in 268 parasite stadia. The lower success rate of the latter may be due to the deteriorating DNA quality since the ITS PCR was performed 5 years after the *cox1* PCR. The ITS rDNA profile was successfully confirmed with Sanger sequencing in a test panel of 12 miracidia. Despite the relatively low number of ITS profiles ($n = 268$), we achieved a good geographical coverage for both markers (Table 1). Table 1 shows the frequency of parasite stadia with a *S. bovis* mitochondrial *cox1* profile for each geographic region and village, per year of sampling.

Averaged over all villages, 79% of the schistosome specimens had a *S. haematobium* (*S.h*) *cox1* profile and 12% had a *S. bovis* (*S.b*) profile (Table 2). The percentage of the *S.b* profile was higher in eggs (19%) than in miracidia (11%). Six percent of the samples displayed a double-banded *Sh/Sb cox1* profile; the majority in eggs (13% compared with 5% in miracidia).

Of the 268 parasites successfully genotyped with ITS rDNA, 243 showed a *S. haematobium* profile (97.9%, Table 2). Three miracidia (1%) presented a hybrid *S.h /S.b* ITS genotype with RFLP, two of them in the village of Pakh and one in Kodit. The hybrid ITS genotype was confirmed in each case by Sanger sequencing; double chromatogram peaks were clearly visible at the diagnostic point mutations between *S. haematobium* (Acc. no FJ588857) and *S. bovis* (Acc. no FJ588862). Two of those miracidia

Table 2. The *cox1* haplotype (A) and ITS rDNA genotype (B) of all schistosome adult worms, eggs and miracidia sampled in Northern Senegal between 2006 and 2010. *S.h/S.b* and *S.h/S.m* indicate that two species bands were observed during the RD-PCR or fragment analysis, or that two peaks were visible after Sanger sequencing.

A)	<i>cox1</i> mtDNA	Adult	Eggs	Miracidia	Total
	<i>S.h</i>	11 (100%)	130 (64.4%)	834 (81.5%)	975 (78.8%)
	<i>S.b</i>	0	38 (18.8%)	110 (10.8%)	148 (12.0%)
	<i>S.h/S.b</i>	0	26 (12.9%)	51 (5.0%)	77 (6.2%)
	<i>S.h/S.m</i>	0	2 (1.0%)	8 (0.8%)	10 (0.8%)
	<i>S.m</i>	0	5 (2.5%)	20 (2.0%)	25 (2.0%)
	Total	11 (0.9%)	202 (16.3%)	1023 (82.8%)	1236 (100%)
B)	ITS rDNA	Adult	Eggs	Miracidia	Total
	<i>S.h</i>	9 (100%)	10 (100%)	224 (97.8%)	243 (97.9%)
	<i>S.h/S.b</i>	0 (0%)	0 (0%)	3 (1.3%)	3 (1.2%)
	<i>S.h/S.m</i>	0 (0%)	0 (0%)	1 (0.4%)	1 (0.4%)
	<i>S.m</i>	0 (0%)	0 (0%)	1 (0.4%)	1 (0.4%)
		9 (3.6%)	10 (4.0%)	249 (92.3%)	268 (100%)

S.h, *S. haematobium*; *S.b*, *S. bovis*; *S.m*, *S. mansoni*.

had a *S.h cox1* profile, while the third was not successfully amplified for *cox1*. Not a single *S.b* ITS profile was found. For 213 parasites we successfully genotyped both *cox1* and ITS rDNA. Twenty of these (9%) showed a *S.b* or a *S.h/S.b cox1* profile with a *S.h* ITS profile and were thus confirmed to be hybrids. This percentage is slightly lower than the total number of hybrids found when adopting the barcoding approach by only taking the *cox1* data into account (12%). The latter statistic is based on a total of 1236 parasite stages (Table 2).

Five eggs and 20 miracidia showed a *S. mansoni cox1* haplotype; their ITS rDNA was not successfully genotyped. We assume that they are pure *S. mansoni* eggs excreted via urine. Two eggs (from Ndiemeul) and eight miracidia (one from Mbodienne, five from Rhonne and two from Ndiemeul) had both an *S.h* and *S.m* band at *cox1*. From five of these the ITS locus was successfully genotyped as *S.h*. In one miracidium the ITS genotype was confirmed to be *S.m* with Sanger sequencing (one insertion in comparison with Acc. no JQ289742) while a double *S.h* and a *S.b* band was found at *cox1*.

Prevalence of *Schistosoma haematobium* and *S. mansoni* infection in humans

The prevalence of *S. haematobium* was high in all regions (Table 2) and ranged between 57.1% in Diatar and 100% in Kodit. The prevalence of *S. mansoni* in stools was highly variable between regions. In the Middle Valley of the Senegal River *S. mansoni* prevalence was lowest with values between 8% (Diatar) and 33% (Dioundiou in 2006). The highest prevalence was found in Rhonne (81%) and in the villages bordering Lac de Guiers,

except for Pakh (17%) (Table 1). The same pattern was observed for co-infection intensities with *S. haematobium* and *S. mansoni* (data not shown).

Geographical distribution of hybrids

The frequency of hybrids, defined as *Schistosoma* stadia with a *S. bovis cox1* profile, was not homogeneously distributed among the four regions in Northern Senegal (Kruskal–Wallis test, $P=0.01$). Villages bordering Lac de Guiers harboured significantly more schistosomes with a *S. bovis cox1* profile (on average 27.5% in 2006 and 2007 and 25.7% in 2009 and 2010) than villages in the Middle Valley of the Senegal River (on average 2.4%, P value Dunn's = 0.01) and the Lampsar River (on average 1.8%, P value Dunn's = 0.046). In Rhonne, in the Delta of the Senegal River, the hybrid frequency was 17% (see Table 2, Fig. 1).

Association between the frequency of hybrids and the prevalence of *Schistosoma* species

The frequency of the *Schistosoma bovis cox1* profile was very high in Diokhor Tack; 74% of the parasite larvae were identified as hybrids. We excluded this outlying observation in the statistical analysis because it is based on parasite eggs from a single host. The PCA biplot (Fig. 2) shows the clustering of villages per geographical region. All villages from the Middle Valley of the Senegal River and the Lampsar River were found at the left side of the first principal component. They had low hybrid frequencies and a relatively low *S. mansoni* prevalence. However, these villages varied in the direction of the second principal component, which corresponds with *S. haematobium* prevalence.




Fig. 2. Biplot of a principal component analysis on the percentage of larval schistosomes with *S. bovis cox1* mtDNA (hybrids), the prevalence of *S. haematobium* (Prev_S.haem) and the prevalence of *S. mansoni* (Prev_S.mans) in nine villages in Northern Senegal by year of sampling. The cumulative proportion is 0.91 for the first two principal components (PC1: 0.61; PC2: 0.30). Observations were coloured per geographical region (Middle Valley Senegal River: yellow; Delta Senegal River: red; Lampsar River: green; Lac de Guiers: colourless). The prevalence of *S. haematobium* and *S. mansoni* was based on egg counts (microscopy), the frequency of hybrids was based on the molecular identification of eggs or miracidia.

Rhonne and the villages bordering Lac de Guiers were found at the right site of the PCA biplot, corresponding to a high frequency of hybrids and a high prevalence of *S. mansoni*. Pakh and Ndieumeul (sampling year 2009/2010) were found at the bottom of the figure. They had the lowest *S. haematobium* prevalence. The incorporation of adults instead of only school-age children (which is the case in sampling year 2009/2010) is expected to lead to a lower *S. haematobium* prevalence. *Schistosoma haematobium* and *S. mansoni* prevalences show a typical age-related pattern with a rapid increase in children, a peak in adolescents and a decline in adults (Meurs et al. 2012). The percentage of hybrids was significantly positively associated with *S. mansoni* prevalence (linear regression: $P = 0.004$, $R^2 = 46.7\%$). Both variables were highest in Rhonne and the villages bordering Lac de Guiers.

Villages that were sampled more than once clustered together, which indicates that over time the frequency of hybrids and the prevalence of *Schistosoma* species remained constant. Only the two observations of the village Mbodienne did not

cluster together: among the 28 genotyped parasites two were hybrids (7%) in 2006, while no hybrids were found among 104 parasites in 2007. The prevalence of *S. haematobium* and *S. mansoni* was also lower in 2007 in Mbodienne.

DISCUSSION

This study supports previous research and confirms that, in Senegal, the majority of hybrid parasites typically present a *Schistosoma haematobium* nuclear (ITS rDNA) and a *S. bovis* mitochondrial (*cox1*) profile (Huyse et al. 2009 and Webster et al. 2013a, b). No pure *S. bovis* ITS profiles were found; only 1% of the ITS profiles displayed both parental profiles. This stands however in contrast with the observation by Webster et al. (2013a, b) in the village Nder at Lac de Guière where 18% ($n = 59$) of the genotyped miracidia from children's urine ($n = 372$) presented a hybrid profile. Of those hybrids, 19% had a *S. haematobium* ITS and *S. bovis cox1* profile while 80% had a *S. bovis* or a double-banded ITS rDNA profile. The low quality of the ITS rDNA PCR in this study might have resulted in an underestimation of *S. bovis* rDNA ITS fragments. However, the difference with the study in Nder may also be due to genuine geographical variation in the distribution of the different types of hybrids.

Barcoding and hybrids

Before discussing the spatiotemporal distribution of hybrids we have to consider the tools used to detect hybrids. Hybrid schistosomes are typically defined based on the discordance between the genetic profile of the mitochondrial *cox1* marker and the nuclear ITS rDNA marker (Sang et al. 1995; Steinauer et al. 2008; Boon et al. 2017). The observed discordance may be the result from historical hybridization followed by introgression or from contemporary hybridization (Detwiler and Criscione, 2010; Hedrick, 2013). In case of the latter, a mixed nuclear profile, indicating a F1 hybrid, would be expected. In case of the former, the ITS profile will have become identical to one of the parental species due to concerted evolution in the ribosomal RNA cistron (Sang et al. 1995). In that case, the hybridization event would be masked and the number of hybrids underestimated. For example, hybrids resulting from a cross between a male *S. bovis* with a female *S. haematobium* may, after a few generations of backcrossing with *S. haematobium*, no longer be detectable with the ITS marker. In the reverse cross, hybridization would still be detectable by the *S. bovis* mitochondrial profile.

When detecting a *S. bovis* mitochondrial profile in schistosome larvae collected from human samples,

one can be absolutely sure of past or ongoing hybridization. Therefore, we argue that the rapid diagnostic PCR assay based on the barcoding fragment is a useful tool to screen large number of human samples, even if it will underestimate the number of hybrids, as it cannot detect the reverse crosses that display a *S. haematobium* *cox1* profile and a *S. bovis* ITS profile. In our study this cross was much less prevalent than the former cross; hybrid parasites with a *S. bovis* ITS fragment must be rare or very focal, as they have been mostly found in one village (Nder) by Webster *et al.* (2013b). When screening cercariae, both the nuclear and the mitochondrial marker should be used, as pure *S. bovis* and pure *S. haematobium* species can sometimes use the same snail species.

Two factors may contribute to the more commonly observed mitochondrial introgression (18%) compared with nuclear introgression (1%): drift and selection (see also below). Because mtDNA is maternally inherited, it is more prone to drift (and indirectly to introgression) in comparison with biparentally inherited nuclear DNA (Ballard and Whitlock, 2004; Currat *et al.* 2008). For example, hybrids have been detected with a complete replacement of mtDNA of one species by another (mitochondrial capture), but without evidence for introgression in the nuclear genome. This pattern was for instance found for two chipmunk species (Good *et al.* 2015). Looking at mitochondrial DNA only may, in this way, provide a biased view of the overall contribution of introgression to the genomes of hybridizing species (Good *et al.* 2015).

Spatiotemporal distribution of hybrids in the SRB

The frequency of hybrid schistosomes remained stable in almost all villages that were sampled twice or more (e.g. for twice in Dioundiou, Mbane; four times in Ndieumeul). Although this study spans only a short time period, it suggests that hybridization between *S. haematobium* and *S. bovis* may already have happened in the past and has led to the spread of, and a stable frequency of *S. bovis* mtDNA among the human population. Hybrid frequency differed between regions, varying between <5% in the Middle Valley of the Senegal River and >30% in villages bordering Lac de Guiers. To explore the potential influence of hybridization in the SRB on schistosomiasis epidemiology, we associated the frequency of hybrids with the prevalence of schistosomiasis. In the discussion below, to find drivers for this heterogeneous pattern, we integrated also data from the literature (e.g. distribution of cattle and, intermediate snail host) with the occurrence of hybrids across the SRB. We start from two hypotheses to explain the geographical heterogeneity. First, in regions with a high hybrid frequency, local conditions may facilitate

hybridization between *S. haematobium* and *S. bovis* (presently or in the past) more than in other regions. Secondly, in regions with a higher frequency of hybrids, the introgression of *S. bovis* mtDNA (or nuclear DNA) may be more adaptive and under positive selection (Irwin, 2002; Toews and Brelsford, 2012).

Link between hybrid distribution and the prevalence of schistosomiasis

The prevalence of *Schistosoma haematobium* was high in all the villages while the prevalence of *S. mansoni* was more patchy with a high prevalence in villages along Lac de Guiers and Rhonne, and a low prevalence in the east of the SRB. The frequency of hybrids was not linked with the prevalence of urinary schistosomiasis. This could advocate against the hypothesis that hybrids would alter schistosomiasis epidemiology, by increasing the transmission of urinary schistosomiasis in the SRB. However, *S. haematobium* prevalence showed little variation across the villages, which makes it less plausible to find a link.

A remarkable observation was the positive association between the frequency of *S. haematobium* × *S. bovis* hybrids and the prevalence of *S. mansoni* parasites. This suggests that in villages with a high *S. mansoni* prevalence there is also a higher risk of infection with hybrids than in villages with mainly urinary schistosomiasis. We slightly violated the conditions for this linear regression: the errors of this model are not fully independent. Some of the villages were sampled twice and there may be some geographical auto-correlation. We did not have sufficient observations to the model dependence of the residuals. However, also within geographical regions and even within villages that were sampled twice we always found the same trend: a positive association between the percentage of hybrids and *S. mansoni* prevalence (Fig. 3).

To explain this association we extrapolate our results (at the village level) to the individual level and look at parasite transmission between individual hosts. Parasites are known to have a different transmission potential in individual hosts with mixed infections in comparison with hosts with single infections (Alizon *et al.* 2013). These interactions can be direct (e.g. competition for host resources, heterologous mating (Tchuem Tchuente *et al.* 1996)) or indirect, by evoking cross-reactive immune responses (Alizon *et al.* 2013). One possible explanation is that once an individual is infected with a schistosome species (e.g. *S. mansoni*), (s)he will become more susceptible to infection by schistosome species that are usually unable to infect humans (e.g. *S. bovis* or hybrid species). For example, *S. mansoni* actively suppresses immune responses (McSorley and Maizels, 2012). In this




Fig. 3. Scatter plot of the association between the percentage of hybrids and *Schistosoma mansoni* prevalence. The linear regression of the percentage of hybrids per village and per year, for all regions combined, is indicated by the dotted regression line ($P = 0.004$, $R^2 = 46.7\%$, regression coefficient = 0.32). The coloured regression lines show the associations for each geographical region separately.

way, *S. mansoni* infections, or mixed infections with *S. mansoni* and *S. haematobium*, may first facilitate infections with *S. bovis* or first generation hybrids, followed by hybridization with *S. haematobium* and ultimately drive the transmission of hybrid schistosomes. In order to test this hypothesis, future studies should focus on the individual level, and link the presence of hybrids within a single host with *Schistosoma* infection intensity. Also, other host-related factors (e.g. host-immunology, ethnicity, water contact behavior) may be associated with the frequency of hybrids. In mycobacteria for example, such variable host-parasite interactions were found, where specific mycobacterial lineages were adapted to particular human populations (Gagneux et al. 2006).

Other explanations of the geographically heterogeneous hybrid distribution

Many other factors may explain or add to the heterogeneous distribution of *S. haematobium* × *S. bovis* hybrids. The occurrence of hybrids may be associated with the (historical) prevalence of *S. bovis*.

Schistosoma bovis inhabits the intestinal veins of cattle, sheep and goats in parts of West Africa including the SRB. A study by Diaw et al. (1998) mentions that along the shores of Lac de Guiers, where most hybrids are found, the breeding of livestock is much more developed and more intensive than in the delta of the Senegal River. As for human schistosomiasis, Diaw et al. (1998) showed also a sharp increase of bovine schistosomiasis and the appearance of new infection foci at the Senegal River Delta and Lac de Guiers after the construction of the dams. Diaw et al. (1998) found that *S. bovis* was more prevalent than *S. curassoni* in Northern Senegal. They related this to the wide distribution of the snail *Bulinus truncatus*, intermediate host of *S. bovis* in Senegal. In a more recent study, no *S. curassoni* was found in the SRB (Webster et al. 2013b); nor were worms with a pure or a hybrid *S. haematobium* profile found in cattle. Hence it remains unknown whether they are a reservoir for hybrid schistosomes. Webster et al. (2013a, b) observed hybrid crosses between *S. haematobium* and *S. curassoni* in children living in Tambacounda and the Vallée du Ferlo but not in the SRB. In the

same regions where *S. haematobium* × *S. curassoni* hybrids were found in children, *S. curassoni* was found in domestic animals. These observations support the hypothesis that the occurrence of schistosome hybrids in humans may be associated with the occurrence of the paternal bovine species in live-stock in the same region. More animal studies are needed to confirm this for the SRB.

Another explanation may be that the introgressed mtDNA is under selection. Mitochondrial DNA has been shown to be highly adaptive (Ballard and Whitlock, 2004; Ballard and Melvin, 2010; Hill, 2015) and important in determining aspects of life-history traits (Ballard and Melvin, 2010). Mitochondria are responsible for the majority of ATP production; 13 genes encode for subunits of the electron transport chain where carbohydrates and fats are oxidized to generate carbon dioxide, water and ATP. The additional genes encode for the translational machinery of the mtDNA itself (Ballard and Whitlock, 2004). It is predicted that important ecological differences are mediated by mitochondrial changes. For example, evolutionary changes in metabolic rates under stress could be effected by changes in the mtDNA sequence (Ballard and Whitlock, 2004). It was found that negative frequency-dependent selection may act on mtDNA and maintain mitochondrial genetic variation (Kazancıoğlu and Arnqvist, 2014). If different mitochondrial haplotypes are favored in different locations, phylogeographic breaks (the occurrence of sharp breaks in mtDNA genealogy that often distinguish regional sets of populations across a species' range) can persist, even in the face of gene flow (Irwin, 2002).

In our setting, one of the two mtDNA *cox1* variants (the *S. bovis* or the *S. haematobium* profile) might be adaptive in certain water bodies or villages, leading to the fixation of the *S. bovis* variant in those areas, and thereby resulting in the observed heterogeneous distribution of hybrids.

Also nuclear *S. bovis* genes may have introgressed that are adaptive or under selection. Several studies have demonstrated evidence for positive selection on introgressed alleles in animals (Harrison and Larson, 2014). For example, a mutation that causes melanism in North American wolves was derived from past hybridization with domestic dogs. The nuclear locus seemed under positive selection in specific habitats: in forested habitats it had risen to high frequencies (Anderson *et al.* 2009). The susceptibility of a snail host depends at least partly on the geographic origin of the snail and the parasite strain (Theron *et al.* 2014). For example, in the Middle Valley (where we found a much lower hybrid frequency), *S. haematobium* is compatible with *B. truncatus* while in the other regions of the SRB *B. globosus* is the main intermediate host (Southgate *et al.* 2000; Sène *et al.* 2004). In light of

this, one may hypothesize that hybrids may have a higher compatibility with the snail strains present around Lac de Guiers compared with other regions in the SRB. Also, the near-total absence of hybrids in the Middle Valley might be linked with the (near-total) absence of *B. globosus* in that area (Huyse, personal observations), if indeed *B. globosus* is a better host for hybrids.

In conclusion, in order to improve insight into potential drivers of hybridization in the SRB, more information on the snail host distribution and the occurrence of *S. bovis* is needed. It is also necessary to screen potential animal reservoirs for hybrid infection. Snail infectivity experiments on the other hand can shed more light on hybrid-snail compatibility.

SUPPLEMENTARY MATERIAL

The supplementary material for this article can be found at <https://doi.org/10.1017/S0031182018000525>

ACKNOWLEDGEMENTS

We thank the field team of Richard Toll, A. Yague, M. Diop, M. Wade and N. Sy for assisting with the sample collection and microscopic analysis. We thank B. Hellemans and S. Geldof for assisting with the molecular work. We also thank Moustapha Mbow for useful suggestions on an earlier version of this manuscript. A special word of thanks goes to the people of the Senegalese villages for their kind participation in the study.

FINANCIAL SUPPORT

N.B. benefited from a Flemish Interuniversity Council (VLIR) VLADOC fellowship and travel grants from the Research Foundation – Flanders and VLIR. Part of the research was covered by a Research Grant of the Research Foundation—Flanders (contract G.0552.10) and a Krediet to navorsers (contract 1516913N) to TH.

ETHICAL STATEMENT

Scientific material was collected in Senegal between 2006 and 2010 in the frame of the SCHISTOINIR project (www.york.ac.uk/res/schistoinir). For this project, approval was obtained from the review board of the Institute of Tropical Medicine (Antwerp, Belgium), the Ethical Committee of Antwerp University Hospital (Antwerp, Belgium) and 'Le Comité National d'Ethique de la Recherche en Santé' (Dakar, Senegal). For the sample from Diatar (1993) approval was obtained from the NHS-LREC of Imperial College (London, UK). Informed consent was obtained from all participants prior to incorporation in the study. At the end of the study inhabitants of participating villages were treated with praziquantel at a dose of 40 mg kg⁻¹ according to World Health Organization (WHO) (WHO, 2006).

CONFLICTS OF INTEREST

None.

REFERENCES

- Abdellahi, M., Ndir, O. and Niang, S. (2016). Évaluation de la prévalence des bilharzioses auprès des enfants de 5 à 14 ans après plusieurs années de traitement de masse dans le bassin du fleuve Sénégal. *Santé Publique* **28**, 535–540.
- Alizon, S., de Roode, J. C. and Michalakis, Y. (2013). Multiple infections and the evolution of virulence. *Ecology Letters* **16**, 556–567.
- Anderson, T. M., Bridgett, M., Candille, S. I., Musiani, M., Stahler, D. R., Smith, D. W., Padhukasahasram, B., Randi, E., Leonard, J. A., Bustamante, C. D., Ostrander, E. A., Tang, H., Robert, K. and Barsh, G. S. (2009). Molecular and evolutionary history of melanism in north American gray wolves. *Science* **323**, 1339–1343.
- Ballard, J. W. O. and Melvin, R. G. (2010). Linking the mitochondrial genotype to the organismal phenotype: invited review. *Molecular Ecology* **19**, 1523–1539.
- Ballard, J. W. O. and Whitlock, M. C. (2004). The incomplete natural history of mitochondria. *Molecular Ecology* **13**, 729–744.
- Barber, K. E., Mkoji, G. M. and Loker, E. S. (2000). PCR-RFLP analysis of the ITS2 region to identify *Schistosoma haematobium* and *S. bovis* from Kenya. *American Journal of Tropical Medicine and Hygiene* **62**, 434–440.
- Boissier, J., Grech-Angelini, S., Webster, B. L., Allienne, J.-F., Huyse, T., Mas-Coma, S., Toulza, E., Barré-Cardi, H., Rollinson, D., Kincaid-Smith, J., Oleaga, A., Galinier, R., Foata, J., Rognon, A., Berry, A., Mouahid, G., Henneron, R., Moné, H., Noel, H. and Mitta, G. (2016). Outbreak of urogenital schistosomiasis in Corsica (France): an epidemiological case study. *The Lancet Infectious Diseases* **3099**, 1–9.
- Boon, N. A. M., Fannes, W., Rombouts, S., Polman, K., Volckaert, F. A. M. and Huyse, T. (2017). Detecting hybridization in African schistosome species: does egg morphology complement molecular species identification? *Parasitology* **144**, 954–964.
- Bowles, J., Blair, D. and McManus, D. P. (1992). Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. *Molecular and Biochemical Parasitology* **54**, 165–173.
- Brant, S. V. and Loker, E. S. (2013). Discovery-based studies of schistosome diversity stimulate new hypotheses about parasite biology. *Trends in Parasitology* **29**, 449–459.
- Brémond, P., Sellin, B., Sellin, E., Nameoua, B., Labbo, R., Theron, A. and Combes, C. (1993). Arguments for the modification of the genome (introgression) of the human parasite *Schistosoma haematobium* by genes from *S. bovis*, in Niger. *Comptes-Rendus de l'Académie des Sciences.Série III : Sciences de la Vie* **316**, 667–670.
- Brindley, P. J. and Hotez, P. J. (2013). Break out: urogenital schistosomiasis and *Schistosoma haematobium* infection in the post-genomic era. *PLoS Neglected Tropical Diseases* **7**, 2–4.
- Curat, M., Ruedi, M., Petit, R. J. and Excoffier, L. (2008). The hidden side of invasions: massive introgression by local genes. *Evolution* **62**, 1908–1920.
- Detwiler, J. T. and Criscione, C. D. (2010). An infectious topic in reticulate evolution: introgression and hybridization in animal parasites. *Genes* **1**, 102–123.
- Diaw, O., Vassiliades, G., Thiongane, Y., Seye, M., Sanrr, Y. and Diouf, A. (1998). Extension des trématodoses du bétail après la construction des barrages dans le bassin du fleuve Sénégal. *Revue d'élevage et de médecinevétérinaire des pays tropicaux* **51**, 113–120.
- Dinno, A. (2016). Dunn's test of multiple comparisons using rank sums. *Dray, S. and Dufour, A. B.* (2007). The ade4 package: implementing the duality diagram for ecologists. *Journal of Statistical Software* **22**, 1–20.
- Dunn, O. J. (1964). Multiple comparisons using rank sums. *Technometrics* **6**, 241–252.
- Emery, A. M., Allan, F. E., Rabone, M. E. and Rollinson, D. (2012). Schistosomiasis collection at NHM (SCAN). *Parasites & Vectors* **5**, 185.
- Gagneux, S., DeRiemer, K., Van, T., Kato-Maeda, M., de Jong, B. C., Narayanan, S., Nicol, M., Niemann, S., Kremer, K., Gutierrez, M. C., Hilty, M., Hopewell, P. C. and Small, P. M. (2006). Variable host-pathogen compatibility in *Mycobacterium tuberculosis*. *Proceedings of the National Academy of Sciences* **103**, 2869–2873.
- Good, J. M., Vanderpool, D., Keeble, S. and Bi, K. (2015). Negligible nuclear introgression despite complete mitochondrial capture between two species of chipmunks. *Evolution* **69**, 1961–1972.
- Harrison, R. G. and Larson, E. L. (2014). Hybridization, introgression, and the nature of species boundaries. *Journal of Heredity* **105**, 795–809.
- Hedrick, P. W. (2013). Adaptive introgression in animals: examples and comparison to new mutation and standing variation as sources of adaptive variation. *Molecular Ecology* **22**, 4606–4618.
- Hill, G. E. (2015). Mitonuclear ecology. *Molecular Biology and Evolution* **32**, 1917–1927.
- Huyse, T., Webster, B. L., Geldof, S., Stothard, J. R., Diaw, O. T., Polman, K. and Rollinson, D. (2009). Bidirectional introgressive hybridization between a cattle and human schistosome species. *PLoS Pathogens* **5**, e1000571.
- Huyse, T., Van den Broeck, F., Hellemans, B., Volckaert, F. A. M. and Polman, K. (2013). Hybridisation between the two major African schistosome species of humans. *International Journal for Parasitology* **43**, 687–689.
- Irwin, D. E. (2002). Phylogeographic breaks without geographic barriers to gene flow. *Evolution* **56**, 2383–2394.
- Kane, R. A., Stothard, J. R., Emery, A. M. and Rollinson, D. (2008). Molecular characterization of freshwater snails in the genus *Bulinus*: a role for barcodes? *Parasites & Vectors* **1**, 15.
- Katz, N., Chaves, A. and Pellegrino, J. (1972). A simple device for quantitative stool thick-smear technique in *Schistosoma mansoni*. *Revista do Instituto de Medicina Tropical de São Paulo* **14**, 397–400.
- Kazancıoğlu, E. and Arnqvist, G. (2014). The maintenance of mitochondrial genetic variation by negative frequency-dependent selection. *Ecology Letters* **17**, 22–27.
- Kruskal, W. H. and Wallis, W. A. (1952). Use of ranks in one-criterion variance analysis. *Journal of the American Statistical Association* **47**, 583–621.
- Leger, E. and Webster, J. P. (2016). Hybridizations within the genus *Schistosoma*: implications for evolution, epidemiology and control. *Parasitology* **144**, 1–16.
- Loker, E. S. and Brant, S. V. (2006). Diversification, dioecy and dimorphism in schistosomes. *Trends in Parasitology* **22**, 521–528.
- McSorley, H. J. and Maizels, R. M. (2012). Helminth infections and host immune regulation. *Clinical Microbiology Reviews* **25**, 585–608.
- Meurs, L., Mbow, M., Vereecken, K., Menten, J., Mboup, S. and Polman, K. (2012). Epidemiology of mixed *Schistosoma mansoni* and *Schistosoma haematobium* infections in northern Senegal. *International Journal for Parasitology* **42**, 305–311.
- Moné, H., Holtfreter, M. C., Allienne, J.-F., Mintsu-Nguéma, R., Ibikounlé, M., Boissier, J., Berry, A., Mitta, G., Richter, J. and Mouahid, G. (2015). Introgressive hybridizations of *Schistosoma haematobium* by *Schistosoma bovis* at the origin of the first case report of schistosomiasis in Corsica (France, Europe). *Parasitology Research* **114**, 4127–4133.
- Morgan, J. A. T., Dejong, R. J., Adeoye, G. O., Ansa, E. D. O., Barbosa, C. S., Brémond, P., Cesari, I. M., Charbonnel, N., Corrêa, L. R., Coulibaly, G., D'Andrea, P. S., De Souza, C. P., Doenhoff, M. J., File, S., Idris, M. A., Incani, R. N., Jarne, P., Karanja, D. M. S., Kazibwe, F., Kpikpi, J., Lwambo, N. J. S., Mabaye, A., Magalhães, L. A., Makundi, A., Moné, H., Mouahid, G., Muchemi, G. M., Mungai, B. N., Sène, M., Southgate, V. et al. (2005). Origin and diversification of the human parasite *Schistosoma mansoni*. *Molecular Ecology* **14**, 3889–3902.
- Norton, A. J., Webster, J. P., Kane, R. A. and Rollinson, D. (2008). Inter-specific parasite competition: mixed infections of *Schistosoma mansoni* and *S. rodhaini* in the definitive host. *Parasitology* **135**, 473–484.
- Picquet, M., Ernould, J. C., Vercruyse, J., Southgate, V. R., Mbaye, A., Sambou, B., Niang, M. and Rollinson, D. (1996). The epidemiology of human schistosomiasis in the Senegal river basin. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **90**, 340–346.
- R Development Core Team (2015). R: A language and environment for statistical computing. R Foundation for Statistical Computing Vienna Austria. doi: 10.1038/sj.hdy.6800737.
- Sang, T., Crawford, D. J. and Stuessy, T. F. (1995). Documentation of reticulate evolution in peonies (*Paeonia*) using internal transcribed spacer sequences of nuclear ribosomal DNA: implications for biogeography and concerted evolution. *Proceedings of the National Academy of Sciences of the United States of America* **92**, 6813–6817.
- Sène, M., Southgate, V. R. and Vercruyse, J. (2004). *Bulinus truncatus*, intermediate host of *Schistosoma haematobium* in the Senegal river basin. *Bulletin de la Société de Pathologie Exotique* **97**, 29–32.
- Soentjens, P., Cnops, L., Huyse, T., Yansouni, C., De Vos, D., Bottieau, E., Clerinx, J. and Van Esbroeck, M. (2016). Diagnosis and clinical management of *Schistosoma haematobium* – *Schistosoma bovis* hybrid infection in a cluster of travelers returning from Mali. *Clinical Infectious Diseases* **63**, 1626–1629.
- Southgate, V., de Clercq, D., Sène, M., Rollinson, D., LY, A. and Vercruyse, J. (2000). Observations on compatibility between *Bulinus*

- truncatus* and *Schistosoma haematobium* in the Senegal river basin. *Annals of Tropical Medicine And Parasitology* **91**, 371–378.
- Southgate, V., Tchuem Tchuente, L., Sène, M., De Clercq, D., Théron, A., Jourdane, J., Webster, B. L., Rollinson, D., Gryseels, B. and Vercruysse, J.** (2001). Studies on the biology of schistosomiasis with emphasis on the Senegal river basin. *Memórias do Instituto Oswaldo Cruz* **96** (Suppl.), 75–78.
- Southgate, V. R., Jourdane, J. and Tchuem Tchuente, L.** (1998). Recent studies on the reproductive biology of the schistosomes and their relevance to speciation in the Digenea. *International Journal for Parasitology* **28**, 1159–1172.
- Steinauer, M. L., Hanelt, B., Mwangi, I. N., Maina, G. M., Agola, L. E., Kinuthia, J. M., Mutuku, M. W., Mungai, B. N., Wilson, W. D., Mkoji, G. M. and Loker, E. S.** (2008). Introgressive hybridization of human and rodent schistosome parasites in western Kenya. *Molecular Ecology* **17**, 5062–5074.
- Tchuem Tchuente, L. A., Southgate, V. R., Combes, C. and Jourdane, J.** (1996). Mating behaviour in schistosomes: are paired worms always faithful? *Parasitology Today* **12**, 231–236.
- Theron, A., Rognon, A., Gourbal, B. and Mitta, G.** (2014). Multi-parasite host susceptibility and multi-host parasite infectivity: a new approach of the *Biomphalaria glabrata*/*Schistosoma mansoni* compatibility polymorphism. *Infection, Genetics and Evolution* **26**, 80–88.
- Toews, D. P. L. and Brelsford, A.** (2012). The biogeography of mitochondrial and nuclear discordance in animals. *Molecular Ecology* **21**, 3907–3930.
- Van den Broeck, F., Geldof, S., Polman, K., Volckaert, F. A. M. and Huyse, T.** (2011). Optimal sample storage and extraction protocols for reliable multilocus genotyping of the human parasite *Schistosoma mansoni*. *Infection, Genetics and Evolution* **11**, 1413–1418.
- Vercruysse, J., Southgate, V. R. and Rollinson, D.** (1984). *Schistosoma curassoni* brumpt, 1931 in sheep and goats in Senegal. *Journal of Natural History* **18**, 969–976.
- Webster, B. L. and Southgate, V. R.** (2003). Compatibility of *Schistosoma haematobium*, *S. intercalatum* and their hybrids with *Bulinus truncatus* and *B. forskalii*. *Parasitology* **127**, 231–242.
- Webster, B. L., Southgate, V. R. and Littlewood, D. T. J.** (2006). A revision of the interrelationships of *Schistosoma* including the recently described *Schistosoma guineensis*. *International Journal for Parasitology* **36**, 947–955.
- Webster, B. L., Rollinson, D., Stothard, J. R. and Huyse, T.** (2010). Rapid diagnostic multiplex PCR (RD-PCR) to discriminate *Schistosoma haematobium* and *S. bovis*. *Journal of Helminthology* **84**, 107–114.
- Webster, B. L., Emery, A. M., Webster, J. P., Gouvras, A., Garba, A., Diaw, O., Seye, M. M., Tchuente, L. A. T., Simoonga, C., Mwangi, J., Lange, C., Kariuki, C., Mohammed, K. A., Stothard, J. R. and Rollinson, D.** (2012). Genetic diversity within *Schistosoma haematobium*: DNA barcoding reveals Two distinct groups. *PLoS Neglected Tropical Diseases* **6**, e1882.
- Webster, B. L., Webster, J. P., Gouvras, A. N., Garba, A., Lamine, M. S., Diaw, O. T., Seye, M. M., Tchuem Tchuente, L. A., Simoonga, C., Mubila, L., Mwangi, J. R., Lwambo, N. J. S., Kabatereine, N. B., Lange, C. N., Kariuki, C., Mkoji, G. M., Rollinson, D. and Stothard, J. R.** (2013a). DNA “barcoding” of *Schistosoma mansoni* across sub-Saharan Africa supports substantial within locality diversity and geographical separation of genotypes. *Acta Tropica* **128**, 250–260.
- Webster, B. L., Diaw, O. T., Seye, M. M., Webster, J. P. and Rollinson, D.** (2013b). Introgressive hybridization of *Schistosoma haematobium* group species in Senegal: species barrier break down between ruminant and human schistosomes. *PLoS Neglected Tropical Diseases* **7**, e2110.
- WHO** (2006). Preventive chemotherapy in human helminthiasis – Coordinated use of anthelmintic drugs in control interventions: a manual for health professionals and programme managers. *World Health Organization* **62**.